

Supplemental table 1. Baseline demographics and clinical characteristics in patients with ≥ 3 prior treatment failures

	Placebo N=286	Erenumab 70 mg N=191	Erenumab 140 mg N=190
N (%)^a	98 (34.3)	69 (36.1)	65 (34.2)
Age, years	42.4 (11.5)	42.8 (11.5)	44.1 (11.3)
Female, n (%)^b	72 (73.5)	62 (89.9)	59 (90.8)
Disease duration, years	24.8 (13.2)	24.5 (13.3)	24.6 (11.9)
MMD	18.6 (4.3)	18.9 (4.4)	19.0 (4.7)
Use of MSM, n (%)	90 (91.8)	62 (89.9)	60 (92.3)
Monthly MSMD	12.0 (7.1)	11.0 (7.6)	12.5 (6.1)
Medication Overuse, n (%)	42 (42.9)	30 (43.5)	27 (41.5)

Data are mean (SD) unless stated otherwise. ^a Percentages represent the number of patients in each prior treatment failure subgroup out of the total patients by treatment subgroup; ^b percentages represent categorical variables compared to the total number of patients in each treatment subgroup by prior treatment failure.

MMD, monthly migraine days; MSM, migraine-specific medication; MSMD, acute migraine-specific medication days; N, number of patients in the analysis set; SD, standard deviation