

Appendix 1

Genetic diversity of dominant plant species in tropical land-use systems on Sumatra, Indonesia

Natalie Breidenbach^{1*}, Sri Rahayu², Iskandar Z. Siregar³, Ulfah J. Siregar³, Ir. Hamzah⁴, Reiner Finkeldey^{1,5}

¹ Georg August University of Göttingen, Department of Forest Genetics and Forest Tree Breeding, Büsgenweg 2, 37077 Göttingen, Germany

² Centre for Plant Conservation, Bogor Botanic Gardens, Indonesian Institute of Science (LIPI), Jl. Ir. H. Juanda, Bogor 16143, Indonesia


³ Department of Silviculture, Faculty of Forestry, Bogor Agricultural University (IPB), Jl. Raya Dramaga, Bogor 16680, Indonesia

⁴ Forestry Faculty, Jambi University, Jl. Lintas Jambi-Muara Bulian Km15, Jambi 36122, Indonesia

⁵ Kassel University, Mönchbergstrasse 19, 34109 Kassel, Germany

*Corresponding author Email: natalie.breidenbach@forst.uni-goettingen.de

Tropical Conservation Science



Appendix 1: Plot design for all 32 plots EFForTS (Drescher et.al.2016). Framed numbers present the positions where dominant plant species were determined. In the tree-dominated systems the “Bitterlich-Method” was applied, in the plantation systems determining the closest species to the 16 positions.

Bukit Dua Belas														Harapan																	
Forest				Jungle Rubber				Oil palm				Rubber				Forest				Jungle rubber				Oil palm				Rubber			
BF1	BF2	BF3	BF4	BJ2	BJ3	BJ4	BJ5	BO2	BO3	BO4	BO5	BR1	BR2	BR3	BR4	HF1	HF2	HF3	HF4	HJ1	HJ2	HJ3	HJ4	HO1	HO2	HO3	HO4	HR1	HR2	HR3	HR4

Appendix 1a: Overview of plot names per land-use systems and region.
 Colours are indicating diversity levels: dark green presents the plot level (α -level), medium green land-use (β -level) and light green the region (γ -level).