

Supplementary Table 1. LIFEware™ characteristics at baseline between those individuals who reported fatigue and those who did not.							
Characteristic	Fatigued, N=2,714	Not Fatigued, N=2,714	OR (95% CI)	Characteristic	Fatigued, N=2,714	Not Fatigued, N=2,714	OR (95% CI)
Get up				Bladder			
Lots of difficulties - Unable	896 (33.2%)	542 (20.0%)		Moderate - Severe	875 (32.6%)	316 (11.8%)	5.4 (4.6-6.4)*
No – Some difficulties	1805 (66.8%)	2165 (80.0%)	2.6 (2.2-3.0)* 1.00 (Ref)	Mild - Moderate	714 (26.6%)	625 (23.3%)	2.0 (1.8-2.3)*
Stairs				No - Mild	1098 (40.9%)	1747 (65.0%)	1.00 (Ref)
Lots of difficulties - Unable	1333 (49.4%)	822 (30.3%)		Vision			
No – Some difficulties	1365 (50.6%)	1887 (69.7%)	3.1 (2.7-3.6)* 1.00 (Ref)	Moderate - Severe	854 (31.6%)	233 (8.6%)	6.3 (5.3-7.5)*
Stand				Mild – Moderate	759 (28.1%)	653 (24.1%)	2.0 (1.7-2.2)*
Lots of difficulties- Unable	1789 (66.4%)	1069 (39.5%)		No - Mild	1092 (40.4%)	1822 (67.3%)	1.00 (Ref)
No – Some difficulties	906 (33.6%)	1639 (60.5%)	4.5 (4.0-5.2)* 1.00 (Ref)	Loneliness			
Drive				Moderate - Extremely	666 (24.7%)	342 (12.0%)	2.4 (2.1-2.8)*
Lots of difficulties- Unable	607 (22.5%)	415 (15.3%)	1.8 (1.6-2.2)*	No - Mildly	2033 (75.3%)	2382 (88.0%)	1.00 (Ref)
No – Some difficulties	1757 (65.1%)	1991 (73.6%)	1.00 (Ref)	Pessimism			
Not applicable	335 (12.4%)	300 (11.1%)	1.4 (1.1-1.6)*	Moderate – Extremely	1174 (43.5%)	636 (23.6%)	2.5 (2.2-2.8)*
Pain				No - Mildly	1525 (56.5%)	2061 (76.4%)	1.00 (Ref)
Pain	1797 (66.4%)	1127 (41.6%)		Tense			
No	909 (33.6%)	1584 (58.4%)	2.8 (2.5-3.1)* 1.00 (Ref)	Moderate – Extremely	1534 (57.3%)	881 (32.7%)	2.8 (2.5-3.2)*
Pain extent			2.4 (2.0-2.8)*	No – Mildly	1144 (42.7%)	1812 (67.3%)	1.00 (Ref)

Discomforting-Excruciating	1425 (79.2%)	700 (62.4%)	1.00 (Ref)	Panic			
Mild pain	374 (20.8%)	422 (37.6%)		Moderate – Extremely	453 (16.8%)	188 (7.0%)	2.7 (2.3-
Upper Limbs				No - Mildly	2242 (83.2%)	2515 (93.0%)	3.3)* 1.00 (Ref)
Moderate - Severe	434 (16.1%)	157 (5.8%)	4.6 (3.8-5.7)*	Annoyed			
Mild - Moderate	668 (24.8%)	336 (12.4%)	3.0 (2.6-3.5)*	Moderate – Extremely	1417 (52.7%)	711 (26.4%)	3.2 (2.8-
No - Mild	1596 (59.2%)	2211 (81.8%)	1.00 (Ref)	No – Mildly	1274 (47.3%)	1987 (73.6%)	3.6)* 1.00 (Ref)
Lower Limbs				Morbid Mood			
Moderate - Severe	1013 (37.6%)	580 (21.5%)	6.4 (5.3-7.6)*	Moderate – Extremely	667 (24.8%)	316 (11.7%)	2.5 (2.2-
Mild - Moderate	823 (30.6%)	528 (19.6%)	3.9 (3.4-4.6)*	No - Mildly	2027 (75.2%)	2385 (88.3%)	2.9)* 1.00 (Ref)
No - Mild	856 (31.8%)	1588 (58.9%)	1.00 (Ref)	Feelings of Guilt			
Bowel				Moderate- Extremely	671 (24.8%)	323 (11.9%)	2.5 (2.1-
Moderate - Severe	574 (21.3%)	163 (6.0%)	5.1 (4.2-6.1)*	No - Mildly	2032 (75.2%)	2381 (88.1%)	2.8)* 1.00 (Ref)
Mild - Moderate	490 (18.2%)	386 (14.3%)	1.8 (1.5-2.1)*				
No - Mild	1625 (60.4%)	2148 (79.6%)	1.00 (Ref)				

Ref=Reference; Odds ratios and confidence intervals are for logistic regression models comparing baseline patient-reported outcomes by fatigue status, and are adjusted for age at enrollment, disease duration, EDSS at enrollment, DMT use, and MS type at enrollment. P-values are also adjusted for multiple comparisons. *p<0.05.