

<i>Score</i>	<i>+0</i>	<i>+1</i>	<i>+2</i>	<i>+3</i>
Aortic neck				
Length (mm)	>25	15-25	10-15	<10
Diameter (mm)	<24	24-26	26-28	>28
Angle (degrees)	>150	150-135	135-120	<120
Calcification (%)	<25	25-50	>50	
Aortic aneurysm				
Tortuosity	<1.15	1.15-1.33	1.33-1.4	>1.4
Angle (degrees)	160-180	140-159	120-139	<120
Thrombus (%)	0	<25	25-50	>50
Branch vessels	None	1 lumbar, IMA	2 vessels <4mm	2 vessels >4mm
Iliac artery				
Pelvic perfusion	Bilateral IIA	Single IIA	Single IIA >50% stenosis	Bilateral IIA occlusion
Calcification (%)	0	<25	25-50	>50
Diameter (mm)				
Minimum	>10	8-10	7-8	<7
Maximum	<20	20-25	25-30	>30
Angle (degrees)	160-180	121-159	90-120	<90
Length (mm)	>30	20-30	10-20	<10

Supplementary Table 1. Anatomic Severity Grading as defined by Chaikof et al¹¹. IIA, Internal iliac artery; IMA, inferior mesenteric artery