

Source	Sum of Squares	DF	Mean Square	F	P
Time	2657,53	5	531,50	13,13	<0,001
Phacic 1 ; Pseudophacic 2	172,78	1	172,78	4,27	0,039
Time*Phacic 1 ; Pseudophacic 2	173,65	5	34,73	0,85	0,50
Residual	18332,71	453	40,47		
Time	Phacic 1 ; Pseudophacic 2	n	Mean	Std. Error	95% Confidence interval
A) Baseline	1	80	22,66	0,71	21,26 to 24,06
	2	22	19,13	1,35	16,47 to 21,80
B) 1st Day	1	80	12,61	0,71	11,21 to 14,01
	2	22	12,04	1,35	9,38 to 14,71
C) 1-3 Months	1	75	15,85	0,73	14,40 to 17,29
	2	17	16,47	1,54	13,43 to 19,50
D) 3-6 Months	1	62	16,25	0,80	14,67 to 17,84
	2	16	14,06	1,59	10,93 to 17,18
E) 9-12 Months	1	54	16,00	0,86	14,29 to 17,70
	2	11	14,90	1,91	11,13 to 18,67
F) 12-18 Months	1	21	16,61	1,38	13,89 to 19,34
	2	5	13,00	2,84	7,40 to 18,59

Table 6. Results of 2 Ways analysis of variance between phacic and pseudophacic eyes in relation to time, regardless the microcatheter system used.