

Appendices

SUPPLEMENTARY TABLE A1. PRO RE NATA (PRN) MEDICATION USE OVER TIME

	CBT for nightmares (n=12)	Treatment as usual (n=12)
PRN antipsychotics		
Week 0	1 (8%)	0
Week 4	1 (8%)	1 (8%)
Week 8	1 (8%)	1 (8%)
PRN mood stabilisers		
Week 0	0 (0%)	0 (0%)
Week 4	0 (0%)	0 (0%)
Week 8	0 (0%)	0 (0%)
PRN anxiolytics		
Week 0	2 (17%)	0 (0%)
Week 4	2 (17%)	1 (8%)
Week 8	2 (17%)	2 (17%)
PRN anti-depressants		
Week 0	0 (0%)	0
Week 4	0 (0%)	1 (8%)
Week 8	0 (0%)	1 (8%)
PRN hypnotics		
Week 0	0 (0%)	0 (0%)
Week 4	0 (0%)	0 (0%)
Week 8	0 (0%)	1 (8%)

Data are n (%).

SUPPLEMENTARY TABLE A2. NIGHTMARE LOG SCORES OVER TIME

	CBT for nightmares	n	Treatment as usual	n	Adjusted mean difference (95% CI)	Effect size (d)
Nightmare log total frequency						
Week 0	7.4 (5.8)	11	9.9 (7.9)	12		
Week 4	4.4 (4.8)	9	8.9 (10.7)	8	-2.6 (-5.8; 0.6)	-0.4
Week 8	3.7 (3.4)	10	12.5 (9.4)	6	-4.8 (-8.3; -1.3)	-0.7
Nightmare log average distress						
Week 0	6.6 (2.4)	11	6.4 (1.8)	12		
Week 4	3.9 (2.9)	9	6.3 (1.7)	9	-2.2 (-4.5; 0.3)	-1.0
Week 8	4.4 (2.9)	10	5.9 (1.5)	6	-1.6 (-4.1; 0.9)	-0.8
Nightmare log – total number of awakenings						
Week 0	6.6 (6.1)	11	7.4 (6.6)	12		
Week 4	3.6 (4.9)	8	7.8 (8.6)	9	-3.7 (-7.4; 0.1)	-0.6
Week 8	1.9 (1.7)	9	9.5 (6.7)	6	-3.8 (-8.0; 0.3)	-0.6
Nightmare log – average sleep quality						
Week 0	1.7 (0.9)	11	1.7 (0.7)	12		
Week 4	2.2 (0.9)	9	1.8 (1.1)	9	0.4 (-0.4; 1.2)	0.5
Week 8	2.1 (0.9)	10	1.4 (0.6)	6	0.6 (-0.3; 1.4)	0.7

Data are mean (SD). CI=confidence interval.

SUPPLEMENTARY TABLE A3. DICHOTOMOUS ASSESSMENT OF SUICIDAL IDEATION ACROSS THE TRIAL PERIOD

	Suicidal ideation outcome at week 4 (n=22)				Suicidal ideation outcome at week 8 (n=20)			
	No suicidal ideation		Suicidal ideation		No suicidal ideation		Suicidal ideation	
	CBT for nightmares	Treatmen t as usual	CBT for nightmares	Treatmen t as usual	CBT for nightmares	Treatmen t as usual	CBT for nightmares	Treatmen t as usual
No suicidal ideation at baseline	6 (27%)	3 (14%)	0 (0%)	0 (0%)	6 (30%)	2 (10%)	0 (0%)	1 (5%)
Suicidal ideation at baseline	0 (0%)	2 (9%)	6 (27%)	5 (23%)	0 (0%)	4 (20%)	5 (25%)	2 (10%)

Data are counts (%). No suicidal ideation indicated by BSS score of 0. Suicidal ideation indicated by BSS >0.

