## **Supporting Information**

A NewAurone from the Seeds of *Psoralea Corylifolia* with Their Diacylglycerol Acyltransferase Inhibitory Activity

Bo Ren<sup>a</sup>, Jia-Yue Dong<sup>a</sup>, Ying Liu<sup>b,c\*</sup>

<sup>a</sup>College of Pharmacy, Chengdu University of Traditional Chinese Medicine, Sichuan Province, 610041, China

<sup>b</sup>Department of Pediatrics, West China Second Hospital of Sichuan University, Sichuan Province, 610041, China

<sup>c</sup>Key Laboratory of Birth Defects and Related Diseases of Women and Children (Sichuan University), Ministry of Education, Sichuan Province, 610041, China

E-mail: liuyingdream@163.com

## Figure Legends:

- Fig. 1 <sup>1</sup>H-NMR spectrum of compound 1 (acetone- $d_6$ , 500 MHz)
- **Fig. 2**  $^{13}$ C-NMR spectrum of compound **1** (acetone- $d_6$ , 125 MHz)
- **Fig. 3** HMBC spectrum of compound 1 (acetone- $d_6$ )
- Fig. 4 HRESIMS spectrum of compound 1





Fig.1 <sup>1</sup>H-NMR spectrum of compound 1 (acetone- $d_6$ , 500 MHz)



**Fig.2** <sup>13</sup>C-NMR spectrum of compound 1 (acetone- $d_6$ , 125 MHz)




Fig.3 HMBC spectrum of compound 1 (acetone- $d_6$ )




Fig.4 HREIMS spectrum of compound 1