SUPPLEMENTARY DATA

A brief up-to-date overview of Amaryllidaceae alkaloids: Phytochemical studies of *Narcissus tazetta* subsp. *tazetta* L., collected in Turkey

Çiğdem Karakoyun ^a, Marco Masi^b, Alessio Cimmino^b, Mustafa Ali Önür^a, Nehir Unver Somer^a, Alexander Kornienko^c and Antonio Evidente ^a,*

^aDepartment of Pharmacognosy, Faculty of Pharmacy, Ege University, 35100 İzmir/ Turkey

^bDepartment of Chemical Sciences, University of Naples Federico II, Complesso Universitario Monte San'Angelo, Via Cintia 4, 80126, Naples, Italy

^cDepartment of Chemistry and Biochemistry, Texas State University, San Marcos, TX 78666, United States

evidente@unina.it

Figure S1. ESI MS spectrum of lycorine (1) recorded in positive modality.




Figure S2. ¹H NMR spectrum of lycorine (1) (CD₃OD on 400 MHz).




Figure S3. ESI MS spectrum of pseudolycorine (21) recorded in positive modality.




Figure S4. ¹H NMR spectrum of pseudolycorine (21) (CD₃OD on 500 MHz).




Figure S5. ESI MS spectrum of galanthamine (22) recorded in positive modality.




Figure S6. ¹H NMR spectrum of galanthamine (**22**) (CD₃OD on 400 MHz).




Figure S7. ESI MS spectrum of 11-hydroxygalanthine (23) recorded in positive modality.




Figure S8. ¹H NMR spectrum of 11-hydroxygalanthine (23) (CD₃OD on 400 MHz).




Figure S9. ¹³C NMR spectrum of 11-hydroxygalanthine (**23**) (CD₃OD on 100 MHz).

