

Literature search full results

Columns: -

Reference

‘Provider’ of outcomes measurement

1. If used only as part of research – Res
2. If used as part of clinical work – Cli
3. If data from ROM - ROM

Recipient of the service

1. Administrators – Adm
2. Clinicians – cli

Burgess objective category

1. To guide clinical decision-making;
2. to engage consumers in treatment;
3. to foster a collaborative approach to care planning and goal setting;
4. to review consumers’ progress with treatment;
5. to inform questions about consumers’ eligibility for given programs;
6. to assist with discharge planning;
7. to improve the evidence base on which services are founded;
8. to evaluate models of service delivery.
9. None of these

Whether cost of using HoNOS was assessed (Y or N)

Whether effectiveness of using HoNOS was assessed (Y or N)

Article reference	Provider	Brief description of purpose of study	recipient	Burgess	cost	effect
Abuelroos T. Factors associated with health and social functioning outcomes following the assessment of a specialist mental health service for adults with developmental disabilities. <i>EmbaseAustralian and New Zealand Journal of Psychiatry</i> . Conference: Royal Australian and New Zealand College of Psychiatrists, ental disabilities. RANZCP Annual Congress 2018. New Zealand. 52 (1 Supplement 1) (pp 75), 2018. Date of Publication: April 2018.	ROM	Supportive evidence in evaluation of a dual disability service	Adm cli	8	N	N
Abuelroos T. Can a specialist mental health service for adults with developmental disabilities make a difference?. <i>EmbaseAustralian and New</i>	ROM	as above	Adm cli	8	N	N

Zealand Journal of Psychiatry. Conference: Royal Australian and New Zealand College of Psychiatrists, RANZCP Annual Congress 2017. Australia. 51 (1 Supplement 1) (pp 33), 2017. Date of Publication: May 2017						
Akande, Emmanuella; Beer, M. Dominic; Ratnajoathy, Kalaanithi. Outcome study of patients exhibiting challenging behaviours four years after discharge from a low secure mental health unit. [References]. <i>PsycINFO Journal of Psychiatric Intensive Care</i> . Vol.3(1), 2007, pp. 21-26	Res ? ROM ?	Follow up study	Cli adm	7	N	N
Allen J ¹ , Hamilton A, Nunn R, Crock L, Frecker J, Burk N. J Psychiatr Ment Health Nurs. 2009 Mar;16(2):129-36. doi: 10.1111/j.1365-2850.2008.01340.x. Evaluation of a community-based mental health drug and alcohol nurse in the care of people living with HIV/AIDS.	ROM	Evaluation of community MH nurse in HIV	adm	8	N	N
Amin S ¹ , Singh SP, Croudace T, Jones P, Medley I, Harrison G. Br J Psychiatry. 1999 May;174:399-403. Evaluating the Health of the Nation Outcome Scales. Reliability and validity in a three-year follow-up of first-onset psychosis.	Res And ROM	Validation study and field trial	adm	9	N	N
Ashaye O ¹ , Mathew G, Dhadphale M. Int J Geriatr Psychiatry. 1997 May;12(5):548-52. A comparison of older longstay psychiatric and learning disability inpatients using the Health of the Nation Outcome Scales.	res	HoNOS used to compare different groups of patients on a psychogeriatric ward	Cli adm	7	N	N
Ashaye, Kunle; Seneviratna, Knightley; Shergill, Sukhwinder; Orrell, Martin. Do the Health of the Nation Outcome Scales predict outcome in the elderly mentally ill? A 1-year follow-up study. [References]. <i>PsycINFO Journal of Mental Health</i> . Vol.8(6), 1999, pp. 615-620	Res	Yes, they do	Cli adm	7	N	N
Ashfaq A., Seneviratne T., Pariente C., Pawlby S. Mental health and mother-infant relationship outcomes on a psychiatric mother and baby unit: Before and after treatment. <i>Embase Archives of Women's Mental Health. Conference: International Marce Society for Perinatal Mental Health Biennial Scientific Conference 2014. Swansea United Kingdom. Conference Publication: (var.pagings). 18 (2) (pp 385-386), 2015. Date of Publication: April 2015</i>	Res	Evaluation of a mother-baby unit	Adm cli	8	N	N
Audin K ¹ , Margison FR, Clark JM, Barkham M. Br J Psychiatry. 2001 Jun;178:561-6. Value of HoNOS in assessing patient change in NHS psychotherapy and psychological treatment services.	ROM	Field trial in psychotherapy setting	cli	9	N	N
Bagley H ¹ , Cordingley L, Burns A, Mozley CG, Sutcliffe C, Challis D, Huxley P. J Clin Nurs. 2000 May;9(3):445-50. Recognition of depression by staff in nursing and residential homes.	Res	What the title says	cli	9	N	N
Baker E., Iqbal E., Johnston C., Broadbent M., Shetty H., Stewart R., Howard R.J., Newhouse S.J., Khondoker M., Kiddle S.J., Dobson R.J.B.	ROM	Clinical measure in follow up study	cli	7	N	N

Dementia severity and progression: Identifying patients at risk for faster cognitive decline. <i>EmbaseAlzheimer's and Dementia. Conference: Alzheimer's Association International Conference, AAIC 2017. United Kingdom. 13 (7) (pp P926), 2017. Date of Publication: July 2017</i>						
Baker E., Iqbal E., Broadbent M., Stewart R., Khondoker M., Newhouse S.J., Dobson R.J.B. Electronic records for dementia research: Do behavioural disturbances, anti-hypertensives and antidepressants influence decline trajectories?. <i>EmbaseAlzheimer's and Dementia. Conference: Alzheimer's Association International Conference 2016. Canada. 12 (7 Supplement) (pp P1056-P1057), 2016. Date of Publication: July 2016</i>	ROM	As above	cli	7	N	N
Barbato A ¹ , Agnetti G, D'Avanzo B, Frova M, Guerrini A, Tettamanti M. J Rehabil Res Dev. 2007;44(6):775-83. Outcome of community-based rehabilitation program for people with mental illness who are considered difficult to treat.	Res	Used as clinical measure in a programme	Adm cli	8	N	N
Bartolommei, N; Benedetti, A; Lattanzi, L; Berti, B; Luchini, F; Cassano, G. B. Depressive and anxiety symptoms in schizophrenia. Impact on course of illness and use of antidepressants in a sample of out-patients. [Italian]. Sintomi depressivi e d'ansia nella schizofrenia. Impatto sul decorso e utilizzo degli antidepressivi in un campione di pazienti trattati in un setting territoriale. <i>PsycINFOGiornale Italiano di Psicopatologia / Italian Journal of Psychopathology. Vol.15(3), 2009, pp. 269-281.</i>	Res	HoNOS a clinical measure in the study	cli	9	N	N
Batla A., Bhatia K.P., Fisher L., Joyce E.M., Price G., Edwards M.J. Outcome after inpatient rehabilitation for patients with functional movement disorders. <i>EmbaseMovement Disorders. Conference: 17th International Congress of Parkinson's Disease and Movement Disorders. Sydney, NSW Australia. Conference Publication: (var.pagings). 28 (SUPPL. 1) (pp S426), 2013. Date of Publication: June 2013</i>	Res	Evaluation of programme	Cli adm	8	N	N
Bauer E, Kleine-Budde K, Stegbauer C, Kaufmann-Kolle P, Goetz K, Bestmann B, Szecsenyi J, Bramesfeld A. BMC Psychiatry. 2016 Jul 15;16:242. doi: 10.1186/s12888-016-0945-z. Structures and processes necessary for providing effective home treatment to severely mentally ill persons: a naturalistic study.	Res	Evaluating factors that relate to better outcomes in HTTs.	adm	7	N	N
Beaglehole B, Frampton CM, Boden JM, Mulder RT, Bell CJ. Aust N Z J Psychiatry. 2017 Nov;51(11):1098-1105. doi: 10.1177/0004867417714879. Epub 2017 Jun 22. An evaluation of Health of the Nation Outcome Scales data to inform psychiatric morbidity following the Canterbury earthquakes.	ROM	Using HoNOS to evaluate the impact of the CCch earthquake on patients	Cli adm	7	N	N

Bebbington P ¹ , Brugha T, Hill T, Marsden L, Window S. Br J Psychiatry. 1999 May;174:389-94. Validation of the Health of the Nation Outcome Scales.	ROM	Field trial	Adm cli	9	N	N
Bech P ¹ , Bille J, Schütze T, Søndergaard S, Wiese M, Waarst S. Nord J Psychiatry. 2003;57(4):285-90. Health of the Nation Outcome Scales (HoNOS): Implementability, subscale structure and responsiveness in the daily psychiatric hospital routine over the first 18 months.	res	Field trial of HoNOS	Cli adm	9	N	N
Beer, M. Dominic; Tighe, James; Ratnajothy, Kalaanithi; Masterson, David. Predicting outcome in low secure environments: A case series from one low secure unit. <i>PsycINFOJournal of Psychiatric Intensive Care. Vol.3(2), 2007, pp. 85-92</i>	ROM	Clinical measure in a specific setting	Cli adm	7	N	N
Bekas, Stavros; Michev, Orlin. Payment by results: Validating care cluster allocation in the real world. <i>PsycINFOThe Psychiatrist. Vol.37(11), 2013, pp. 349-355.</i>	ROM	Review of 600 cases of use of clustering tool and finding reasons for mismatches (clinical judgment is final arbiter)	Cli adm	7	N	N
Berdzenishvili E., Chkonia E., Okruashvili M., Roinishvili M. Cognitive deficit and social functioning in patients with first psychotic episode. <i>EmbaseEuropean Neuropsychopharmacology. Conference: 28th European College of Neuropsychopharmacology, ECNP Congress. Amsterdam Netherlands. Conference Publication: (var.pagings). 25 (SUPPL. 2) (pp S332-S333), 2015. Date of Publication: September 2015.</i>	Res	Part of evaluation of FEP patients	cli	9	N	N
Berendsen S, van der Paardt J, van Bruggen M, Nusselder H, Jalink M, Peen J, Van H, Dekker J, de Haan L. Clin Schizophr Relat Psychoses. 2018 Jun 26. doi: 10.3371/CSRP.BEPA.061518. [Epub ahead of print] Exploring construct validity of clinical staging in schizophrenia spectrum disorders in an acute psychiatric ward.	Cli ROM	HoNOS used as a measure in the evaluation of the construct validity of clinical staging in schizophrenia	cli	9	N	N
Richard H., Campbell N., Newby G., Odiyoor M., Pinder C. Dual diagnosis, singular care: A joint neuropsychiatry, neuropsychology and neurorehabilitation clinic for those with acquired brain injury and psychiatric comorbidities. <i>EmbaseBrain Injury. Conference: 11th World Congress on Brain Injury of the International Brain Injury Association. Netherlands. 30 (5-6) (pp 575), 2016. Date of Publication: 2016.</i>	ROM	Evaluation of new model of care	Adm cli	8	N	N
Bjørngaard JH ¹ , Ruud T, Friis S. Soc Psychiatry Psychiatr Epidemiol. 2007 Oct;42(10):803-9. Epub 2007 Jul 11. The impact of mental illness on patient satisfaction with the therapeutic relationship: a multilevel analysis.	Res	HoNOS used as clinical measure in a satisfaction survey	Cli adm	9	N	N
Blair, Grant; Deane, Carl. Inner-city general practice population of people with schizophrenia. <i>PsycINFOThe Psychiatric Bulletin. Vol.22(4), 1998, pp. 221-225</i>	Res	Attempt to describe the MH population in inner city GP	Cli adm	7	N	N

Blankers M., van Emmerik A., Richters B., Dekker J. Blended internet care for patients with severe mental illnesses: An open label prospective controlled cohort pilot study. <i>EmbaseInternet Interventions</i> . 5 (pp 51-55), 2016. Date of Publication: 01 Sep 2016	Res	Assessment of incorporation of internet contact into ACT	cli	8	N	N
Boehnke J.R., Aragon M.J., Moran V., Clark M., Chalkley M., Jacobs R. The mental health clustering tool (MHCT) in NHS England's secondary care: Does the available evidence support a payment by results approach?. <i>EmbaseQuality of Life Research. Conference: 24th Annual Conference of the International Society for Quality of Life Research, ISOQOL 2017. United States</i> . 26 (1 Supplement 1) (pp 2), 2017. Date of Publication: October 2017	ROM	Evaluation of 2 clustering models	adm	8	N	N
Bonsack C., Borgeat F., Lesage A. Measuring patients problems severity and outcomes in a psychiatric sector: A field study with the French version of the Health of Nation Outcome Scales (HoNOS-F). <Mesurer la severite des problemes des patients et leur evolution dans un secteur psychiatrique: Une etude sur le terrain du Health of Nation Outcome Scales en Francais (HoNOS-F).> <i>EmbaseAnnales Medico-Psychologiques</i> . 160 (7) (pp 483-488), 2002. Date of Publication: September 2002	Res	Empirical field trial	Cli Adm	9	N	N
Boot B ¹ , Hall W, Andrews G. Br J Psychiatry. 1997 Sep;171:242-6. Disability, outcome and case-mix in acute psychiatric in-patient units.	Res	Field trial of HoNOS & another with in-patients	Cli adm	9	N	N
Briner D, Jäger M, Kawohl W, Baumgartner-Nietlisbach G. Psychiatr Prax. 2017 Sep;44(6):339-347. doi: 10.1055/s-0042-111742. Epub 2016 Sep 14. [Mental Disorder and Self-Rated Health Among Homeless People in Zurich - First Epidemiological Data from Switzerland]. [Article in German; Abstract available in German from the publisher]	res	Assessment of psychiatric morbidity of homeless people	Cli adm	9	N	N
Brooker C ¹ , Molyneux P, Deverill M, Repper J. J Adv Nurs. 1999 Jan;29(1):44-51. Evaluating clinical outcome and staff morale in a rehabilitation team for people with serious mental health problems.	Res	HoNOS score an independent variable in a study of staff morale	Cli adm	7	N	n
Brooker, Charlie; Molyneux, Philip; Deverill, Mark; Repper, Julie. An audit of costs and outcome using HoNOS-3 in a rehabilitation team: A pilot study. <i>PsycINFOJournal of Mental Health</i> . Vol.6(5), 1997, pp. 491-502.	Res	costs of formal and informal treatment, not of using HoNOS; mostly a field trial	Cli adm	9	N	n
Bruce J, Watson D, van Teijlingen ER, Lawton K, Watson MS, Palin AN. J Adv Nurs. 1999 May;29(5):1060-7. Dedicated psychiatric care within general practice: health outcome and service providers' views.	Res	Study of use of CPN in general practice	Adm cli	8	N	N
Burgess P, Pirkis J, Coombs T. Aust New Zealand Health Policy. 2006 Aug 30;3:9. Do adults in contact with Australia's public sector mental health services get better?	ROM	describes the benefit of treatment, not the benefit of the use of ROM.	adm	7	N	N
Burman-roy S, Butterworth M, Madan I, Henderson M, Harvey SB.	ROM	Justifying an occupational psychiatry service	adm	8	N	N

Occup Med (Lond). 2013 Oct;63(7):507-9. doi: 10.1093/occmed/kqt103. Epub 2013 Aug 29. Which patients are seen by an occupational psychiatry service?						
Carta MG, Agaj A, Harapej E, Lecca ME, Xhelili G, Altoé G, Mura G, Moro MF, Angermeyer MC. Int J Soc Psychiatry. 2013 Nov;59(7):682-9. doi: 10.1177/0020764012452353. Epub 2012 Jul 17. Outcomes of discharged females versus those waiting for discharge from Vlore Psychiatric Hospital (Albania).	ROM	Evaluation of home group discharge compared to continuing hospitalisation	Adm cli	8	N	N
Carta MG ¹ , Angermeyer MC, Sancassiani F, Tuligi F, Pirastu R, Pisano A, Pintus E, Mellino G, Pintus M, Pisanu E, Moro MF, Massidda D, Trincas G, Bhugra D. BMC Psychiatry. 2013 Dec 6;13:333. doi: 10.1186/1471-244X-13-333. A follow-up on patients with severe mental disorders in Sardinia after two changes in regional policies: poor resources still correlate with poor outcomes.	Res	Increasing resources improves outcomes	adm	7	N	N
Carta MG, Maggiani F, Pilutzu L, Moro MF, Mura G, Cadoni F, Sancassiani F, Vellante M, Machado S, Preti A. Clin Pract Epidemiol Ment Health. 2014 Jul 23;10:73-9. doi: 10.2174/1745017901410010073. eCollection 2014. Sailing for rehabilitation of patients with severe mental disorders: results of a cross over randomized controlled trial.	Res	Evaluation of a sailing rehabilitation programme	cli	8	N	N
Carta M ¹ , Sancassiani F ¹ , Lecca M ¹ , Pintus E ¹ , Pintus M ¹ , Pisano E ¹ , Congiu S ¹ , Moro M ¹ , Angermeyer C ² . Clin Pract Epidemiol Ment Health. 2013 Nov 28;9:214-20. doi: 10.2174/1745017901309010214. eCollection 2013. Coping with the crisis: people with severe mental disorders acting for social change through sustainable energy.	Res	Evaluation of a work inclusion programme	Cli adm	8	N	N
Castelein S., Swart M., Wolters G., Bruggeman R., Kneegting H. Does routine outcome monitoring in schizophrenia improve the implementation of evidence based care?. <i>EmbaseSchizophrenia Research. Conference: 4th Biennial Schizophrenia International Research Conference. Florence Italy. Conference Publication: (var.pagings). 153 (SUPPL. 1) (pp S185), 2014. Date of Publication: April 2014</i>	ROM	The answer - not necessarily; good study	Adm cli	7	N	YES
Catts S.V., O'Toole B.I., Carr V.J., Lewin T., Neil A., Harris M.G. Early psychosis intervention: Service model determines case detection rate but what the clinician does determines 6 to 12 month outcomes. <i>EmbaseSchizophrenia Research. Conference: 3rd Biennial Schizophrenia International Research Conference. Florence Italy. Conference Publication: (var.pagings). 136 (SUPPL. 1) (pp S178), 2012. Date of Publication: April 2012</i>	ROM	EIP programmes improve early detection but not outcomes	Cli adm	8	N	N

Chaplin, Robert; Perkins, Rachel. HoNOS : A cautionary tale of their use in a rehabilitation service. <i>PsycINFOThe Psychiatric Bulletin</i> . Vol.23(1), 1999, pp. 20-21	Res	Field trial – poor inter-rater reliability or correspondence with clinical improvement	Cli adm	9	N	N
Cheung G ¹ , Strachan J. N Z Med J. 2007 Aug 10;120(1259):U2660. Routine 'Health of the Nation Outcome Scales for elderly people' (HoNOS65+) collection in an acute psychogeriatric inpatient unit in New Zealand.	ROM	Description of HoNOS score reductions in a psychogeriatric service	adm	7	N	N
Christian-Jones C., Hastings R., Hughes J., Lindsay W., Roberts D. The associations between physical health, mental health and challenging behavior in adults with an intellectual disability living in residential services. <i>EmbaseJournal of Intellectual Disability Research. Conference: 14th World Congress of the International Association for the Scientific Study of Disabilities, IASSID 2012. Halifax, NS Canada. Conference Publication: (var.pagings). 56 (7-8) (pp 687), 2012. Date of Publication: July 2012</i>	Res	Part of a survey of patients	cli	7	N	N
Cocchi A ¹ , Meneghelli A, Preti A. Aust N Z J Psychiatry. 2008 Dec;42(12):1003-12. doi: 10.1080/00048670802512032. Programma 2000: celebrating 10 years of activity of an Italian pilot programme on early intervention in psychosis.	ROM	Description of an EIP service	adm	9	N	N
Cook, Sarah; Howe, Amanda. Engaging people with enduring psychotic conditions in primary mental health care and occupational therapy. <i>PsycINFOThe British Journal of Occupational Therapy</i> . Vol.66(6), 2003, pp. 236-246.	Res	Trial of new model of inreach	Adm ali	8	N	N
Corrivetti G., Caserta D., Perna C. Implementation of cognitive rehabilitation in psychiatric services: Integration of cognitive remediation and psychosocial rehabilitation. <i>EmbaseJournal of Psychopathology</i> . 24 (2) (pp 93-97), 2018. Date of Publication: June 2018	Res	Part of the assessment of a cognitive rehabilitation programme	Cli adm	8	N	N
Crawford MJ ¹ , Zoha M, Macdonald AJD, Kingdon D. BJPsych Bull. 2017 Jun;41(3):172-176. doi: 10.1192/pb.bp.116.054346. Improving the quality of mental health services using patient outcome data: making the most of HoNOS.	ROM	Description of how 3 MHSs have used HoNOS 1 – to examine reasons for admission 2 – identifying which patients do worse in ATTs led to development of more specific programmes for them. 3 – HoNOS identified an old-age unit with declining outcomes in depression; led to a further study, which picked up a link between routine prescribing of benzos and poorer	cli	4	N	YES

		outcomes; changing practice improved outcomes.				
Crockford, Helena A; Brunton, Joan; Surgenor, Tammy E. Clinical effectiveness of an acute psychiatric day hospital run on therapeutic community lines. <i>PsycINFOTherapeutic Communities</i> . Vol.24(1), 2003, pp. 37-54.	ROM	Evaluation of day hospital	Adm cli	8	N	N
Dalal B., Larkin E., Leese M., Taylor P.J. Clozapine treatment of long-standing schizophrenia and serious violence: A two-year follow-up study of the first 50 patients treated with clozapine in Rampton high security hospital. <i>EmbaseCriminal Behaviour and Mental Health</i> . 9 (2) (pp 168-178), 1999. Date of Publication: 1999.	Res	Uncontrolled drug trial	cli	8	N	N
Dalton, Rita. Client progress in a community team for mentally disordered offenders. <i>PsycINFOThe British Journal of Forensic Practice</i> . Vol.7(1), 2005, pp. 18-22	Res	Review of new service	Adm cli	8	N	N
D'Antonio, Minh-Lieu Veronica. Evaluating the effectiveness of specialist multi-disciplinary community teams for people with intellectual disabilities in the United Kingdom. <i>PsycINFODissertation Abstracts International: Section B: The Sciences and Engineering</i> . Vol.66(12-B),2006, pp. 6917	Res	What the title says	Adm cli	8	N	N
D'Avanzo B ¹ , Aliprandini E, Beghi M, Cornaggia CM, Erlicher A, Frova M, Mascarini A, Miragoli P, Righi A. Epidemiol Psichiatri Soc. 2008 Jan-Mar;17(1):57-64. [Residential facilities and day centres in mental health. Is there any difference?]. [Article in Italian]	Res	As a clinical measure in comparing two patient groups in different settings	Adm cli	9	N	N
D'Avanzo B ¹ , Battino RN, Gallus S, Barbato A. Aust N Z J Psychiatry. 2004 Aug;38(8):619-28. Factors predicting discharge of patients from community residential facilities: a longitudinal study from Italy.	res	HoNOS a severity measure in a follow up study	Cli adm	9	N	N
de Beurs E, Blankers M, Delespaul P, van Duijn E, Mulder N ⁷ , Nugter A, Swildens W, Tiemens BG, Theunissen J, van Voorst AFA, van Weeghel J. BMC Psychiatry. 2018 Jul 13;18(1):225. doi: 10.1186/s12888-018-1798-4. Treatment results for severe psychiatric illness: which method is best suited to denote the outcome of mental health care?	adm	Comparing other outcomes measures with HoNOS for ranking 12/12 outcomes in institutions	adm	8	N	N
Deenik J., Tenback D., Hendriksen I., Tak E., Van Harten P. Multidisciplinary lifestyleenhancing treatment for inpatients with severe mental illness (multi-study): Effects on physical health, psychotic symptoms, quality of life and functioning. <i>EmbaseSchizophrenia Bulletin. Conference: 6th Biennial Schizophrenia International Research Society Conference, SIRS 2018. Italy. 44 (Supplement 1) (pp S407), 2018. Date of Publication: April 2018</i>	Res	Evaluation of a new IP programme	Cli adm	8	N	N

de Girolamo G; Candini V; Buizza C; Ferrari C; Boero ME; Giobbio GM; Goldschmidt N; Greppo S; Iozzino L; Maggi P; Melegari A; Pasqualetti P; Rossi G. Is psychiatric residential facility discharge possible and predictable? A multivariate analytical approach applied to a prospective study in Italy. <i>Ovid MEDLINE(R)Social Psychiatry & Psychiatric Epidemiology</i> . 49(1):157-67, 2014 Jan	Res	HoNOS used as an independent variable, looking at discharge prediction	cli	9	N	N
de Girolamo G ¹ , Picardi A, Santone G, Falloon I, Morosini P, Fioritti A, Micciolo R; PROGRES Group. Psychol Med. 2005 Mar;35(3):421-31. The severely mentally ill in residential facilities: a national survey in Italy.	Res	Part of a survey of patients in residential (non-hospital) facilities	Cli adm	7	N	N
Demartini B, Batla A, Petrochilos P, Fisher L, Edwards MJ, Joyce E. J Neurol. 2014 Dec;261(12):2370-7. doi: 10.1007/s00415-014-7495-4. Epub 2014 Sep 20. Multidisciplinary treatment for functional neurological symptoms: a prospective study.	Res	Service evaluation through a follow up study	Cli adm	7	N	N
de Girolamo G ¹ , Candini V, Buizza C, Ferrari C, Boero ME, Giobbio GM, Goldschmidt N, Greppo S, Iozzino L, Maggi P, Melegari A, Pasqualetti P, Rossi G. Soc Psychiatry Psychiatr Epidemiol. 2014 Jan;49(1):157-67. doi: 10.1007/s00127-013-0705-z. Epub 2013 May 28. Is psychiatric residential facility discharge possible and predictable? A multivariate analytical approach applied to a prospective study in Italy.	Res	HoNOS used as an independent variable against likelihood of discharge from a psychiatric residential facility	cli	7	N	N
Den Brok W., Sterkenburg P., Schuengel C. Using mobile technology to support relationship development and emotional well-being: A case study. <i>EmbaseJournal of Intellectual Disability Research. Conference: 14th World Congress of the International Association for the Scientific Study of Disabilities, IASSID 2012. Halifax, NS Canada. Conference Publication: (var.pagings). 56 (7-8) (pp 680), 2012. Date of Publication: July 2012</i>	res	Use of technology supported	cli	8	N	N
De Risio A, Pancheri A, Simonetti G, Giannarelli D, Stefanutto L, Gentile B. Prog Neuropsychopharmacol Biol Psychiatry. 2011 Jun 1;35(4):1112-6. doi: 10.1016/j.pnpbp.2011.03.011. Epub 2011 Apr 5. Add-on of aripiprazole improves outcome in clozapine-resistant schizophrenia.	Res	Used as outcome measure for drug study	cli	9	N	N
Devlin R., Medel N., Jairam R., Barton G., Eapen V. Making sense of chaos - A brave foray into helping the most disadvantaged: A joint mental health, drug health and family and community services initiative. <i>EmbaseAustralian and New Zealand Journal of Psychiatry. Conference: Royal Australian and New Zealand College of Psychiatrists, RANZCP Annual Congress 2018. New Zealand. 52 (1 Supplement 1) (pp 70), 2018. Date of Publication: April 2018</i>	ROM	Evaluation of a new team	Cli adm	8	N	N
Devlin R., Medel N., Jairam R., Barton G., Eapen V. What we have learnt working with the first few families in the whole-family team?.	ROM	Evaluation of a new team	Cli adm	8	N	N

EmbaseAustralian and New Zealand Journal of Psychiatry. Conference: Royal Australian and New Zealand College of Psychiatrists, RANZCP Annual Congress 2018. New Zealand. 52 (1 Supplement 1) (pp 72), 2018. Date of Publication: April 2018.						
Dickens, Geoff; Sugarman, Philip; Picchioni, Marco; Long, Clive. HoNOS-Secure: Tracking risk and recovery for men in secure care. PsycINFOThe British Journal of Forensic Practice. Vol.12(4), 2010, pp. 36-46	Res ? ROM	Field trial of use	adm	7	N	N
Dickens G., Sugarman P. Key performance indicators and HoNOS. Ovid EmcareBritish Journal of Health Care Management. 16 (9) (pp 431-435), 2010. Date of Publication: September 2010	ROM	Use stable HoNOS results findings to advocate for wider use of improvements rates as a KPI	adm	9	N	N
Draper, Brian; Busetto, Giselle; Cullen, Breda. Risk factors for and prediction of falls in an acute aged care psychiatry unit. PsycINFOAustralasian Journal on Ageing. Vol.23(1), 2004, pp. 48-51.	Res	Clinical tool used in a prediction study	cli	9	N	N
Dratcu L., Walker-Tilley T., Ramanuj P., Lopez-Morinigo J., Huish E. Metropolitan acute hospital care in psychiatry: Measuring outcomes. EmbaseEuropean Psychiatry. Conference: 20th European Congress of Psychiatry, EPA 2012. Prague Czech Republic. Conference Publication: (var.pagings). 27 (SUPPL. 1) (no pagination), 2012. Date of Publication: 2012. [Conference Abstract]	ROM	Not sure why they did this study	Adm cli	9	N	N
Drukker M, Visser E, Sytema S, van Os J. Clin Pract Epidemiol Ment Health. 2013 Nov 28;9:202-9. doi: 10.2174/1745017901309010202. eCollection 2013. Flexible Assertive Community Treatment, Severity of Symptoms and Psychiatric Health Service Use, a Real life Observational Study.	ROM	Evaluation of an ACT programme	Cli adm	8	N	N
Dunt DR, Benoy AW, Phillipou A, Collister LL, Crowther EM, Freidin J, Castle DJ. Aust Health Rev. 2017 Oct;41(5):573-581. doi: 10.1071/AH16055. Evaluation of an integrated housing and recovery model for people with severe and persistent mental illnesses: the Doorway program.	Cli ROM	Evaluation of a housing & recovery support programme	adm	8	N	N
Egger ST, Vetter S, Weniger G, Vandeleur C, Seifritz E, Müller M. Front Public Health. 2016 Oct 13;4:220. eCollection 2016. The Use of the Health of the Nation Outcome Scales for Assessing Functional Change in Treatment Outcome Monitoring of Patients with Chronic Schizophrenia.	Res	Use of HoNOS as a measure of function, compared with PANSS as a measure of symptoms. Purpose to encourage more efforts towards functional improvement, not just symptomatic	cli	1	N	N
El Aissati D., Jackson C. Birmingham early intervention service redesign: Improving recovery and social outcomes for young people with psychosis.	ROM	Evaluation of new recovery orientation in eip	cli	8	N	N

<i>EmbaseEarly Intervention in Psychiatry. Conference: 8th International Conference on Early Psychosis: From Neurobiology to Public Policy. San Francisco, CA United States. Conference Publication: (var.pagings). 6 (SUPPL.1) (pp 86), 2012. Date of Publication: October 2012</i>						
El-Badri S ¹ , Mellsop G. <i>Australas Psychiatry</i> . 2011 Oct;19(5):410-4. doi: 10.3109/10398562.2011.602078. Epub 2011 Aug 24. Clozapine use and outcomes among patients with treatment resistant schizophrenia.	ROM	Study of clozapine use and a variety of other variables, including HoNOS	cli	9	N	N
Eskelinen S., Suvisaari J. Do schizophrenic out-patients receive appropriate somatic care?. <i>EmbaseSchizophrenia Research. Conference: 2nd Schizophrenia International Research Society Conference, SIRS 2010. Florence Italy. Conference Publication: (var.pagings). 117 (2-3) (pp 290), 2010. Date of Publication: April 2010</i>	Res	Part of a survey of medical needs	cli	7	N	N
Fankhauser S, Hochstrasser B, Sievers M, Soyka M. <i>Psychother Psychosom Med Psychol</i> . 2017 Sep;67(9-10):391-400. doi: 10.1055/s-0043-105482. Epub 2017 Jun 29. [Assessing Change of Depressive Symptoms and Severity of Depression in an Inpatient Setting : Performance of the HoNOS (Health of the Nation Outcome Scales)]. [Article in German; Abstract available in German from the publisher]	Cli	Assessment of the value of HoNOS in assessing depression – other tools needed	res	9	N	N
Fava S., De Girolamo G., Picardi A., et al. Drug prescription in Italian Residential Facilities. <La prescrizione di psicofarmaci nelle Strutture Residenziali Italiane.> <i>EmbaseEpidemiologia e Psichiatria Sociale</i> . 14 (2) (pp 77-90), 2005. Date of Publication: April/June 2005.	Res	As a clinical measure in a study of prescribing	cli	7	N	N
Fisher LJ, Goldney RD, Furze PF, Williams JL, Mattner J, McCleave DJ. <i>J ECT</i> . 2004 Sep;20(3):174-8. Electroconvulsive therapy, depression, and cognitive outcomes: an Australian audit.	Res	ECT follow up study	cli	9	N	N
Fankhauser, Sonja; Hochstrasser, Barbara; Sievers, Marcus; Soyka, Michael. Assessing change of depressive symptoms and severity of depression in an inpatient setting: Performance of the HoNOS (Health of the Nation Outcome Scales). [German]. Die Eignung der HoNOS (Health of the Nation Outcome Scales) zur Erfassung des Verlaufs und des Schweregrads depressiver Symptomatik im stationären Setting. <i>PsycINFOPPmP: Psychotherapie Psychosomatik Medizinische Psychologie</i> . Vol.67(9-10), 2017, pp. 391-400.	Res	Comparing HoNOS change in different levels of depressive IPs	cli	9	N	N
Ferrara M., Guloksuz S., Burke S., Baccari F., Miselli M., Saponaro A., Ferri M., Srihari V., Starace F., Psicotici G.R.E. Implementation of a program for early intervention in psychosis onset: The experience of regione Emilia Romagna, Northern Italy. <i>EmbaseSchizophrenia Bulletin. Conference: 6th Biennial Schizophrenia International Research Society</i>	Res	Evaluation of early intervention	Cli adm	8	N	N

Conference, SIRS 2018. Italy. 44 (Supplement 1) (pp S426-S427), 2018. Date of Publication: April 2018						
Fisher, Laura J; Goldney, Robert D; Furze, Peter F; Williams, Jody L; Mattner, June; McCleave, David J. Electroconvulsive therapy, depression, and cognitive outcomes. <i>PsycINFOThe Journal of ECT</i> . Vol.20(3), 2004, pp. 174-178	Res	Clinical measure in ect study	cli	8	N	N
Frost BG, Turrell M, Sly KA, Lewin TJ, Conrad AM, Johnston S, Tirupati S, Petrovic K, Rajkumar S. BMC Health Serv Res. 2017 Jan 3;17(1):2. doi: 10.1186/s12913-016-1939-8. Implementation of a recovery-oriented model in a sub-acute Intermediate Stay Mental Health Unit (ISMHU).	ROM	justification of a new unit	adm	8	N	N
Gallagher J, Teesson M. Aust N Z J Psychiatry. 2000 Oct;34(5):850-5. Measuring disability, need and outcome in Australian community mental health services.	ROM	Field trial	Cli adm	9	N	N
Galvan F., Pisano A., Meneghelli A. The goal of social recovery: Therapeutic and rehabilitative groups in an early intervention program. <i>EmbaseEarly Intervention in Psychiatry. Conference: 7th International Conference on Early Psychosis - Early Psychoses: A Lifetime Perspective. Amsterdam Netherlands. Conference Publication: (var.pagings). 4 (SUPPL. 1) (pp 152), 2010. Date of Publication: November 2010.</i>	res	Evaluation of EIP	Cli adm	8	N	N
Ganeshalingam, Yogesh; Chang, David; Dunn, Nick. The evidence for effective psychiatric admissions as measured by an audit of health of the nation outcome scales recording. <i>PsycINFOJournal of Evaluation in Clinical Practice</i> . Vol.16(1), 2010, pp. 196-198	ROM	recommendation for using HoNOS to determine treatment effectiveness	Adm cli	7	N	N
Giacomin M. Does community care work? Evaluate the effectiveness of mental health services. <i>EmbaseEuropean Psychiatry. Conference: 24th European Congress of Psychiatry, EPA 2016. Madrid Spain. Conference Publication: (var.pagings). 33 (SUPPL.) (pp S631), 2016. Date of Publication: March 2016.</i>	Res	Comparison of 3 treatment packages	cli	7	N	N
Giordano V., Paramel A., Gupta S. Community treatment order experience and implications in a socially deprived inner city London borough. <i>EmbaseEuropean Psychiatry. Conference: 21st European Congress of Psychiatry, EPA 2013. Nice France. Conference Publication: (var.pagings). 28 (SUPPL. 1) (no pagination), 2013. Date of Publication: 2013</i>	ROM	Clinical dependent variable in a study of CTO effectiveness	Cli adm	8	N	N
Girardi, Alessandra; Snyman, Pieter; Natarajan, Muthusamy; Griffiths, Christopher. Older adults in secure mental health care: Health, social wellbeing and security needs measured with honos-secure across different age groups. <i>PsycINFOJournal of Forensic Psychiatry & Psychology</i> . 2018, pp. No Pagination Specified.	ROM	Comparison of different ages in a secure MH unit	cli	7	N	N

Goldney RD, Fisher LJ, Walmsley SH. Aust N Z J Psychiatry. 1998 Apr;32(2):199-205. The Health of the Nation Outcome Scales in psychiatric hospitalisation: a multicentre study examining outcome and prediction of length of stay.	res	Field trial of HoNOS , testing for predictability of LOS	Cli adm	7	N	N
Green, Heather J; Frank, Rowena M; Butwell, Andrew; Beck, Oswald J. Implementation and evaluation of brief cognitive-behavioural therapy in a mental health acute assessment and treatment service. <i>PsycINFOBehaviour Change</i> . Vol.24(2), 2007, pp. 87-98.	Res	Evaluation of cbt	Cli Adm	8	N	N
Griffiths C., Roychowdhury A., Girardi A. Seclusion: the association with diagnosis, gender, length of stay and HoNOS-secure in low and medium secure inpatient mental health service. <i>EmbaseJournal of Forensic Psychiatry and Psychology</i> . (pp 1-18), 2018. Date of Publication: 14 Feb 2018	ROM	HoNOS probably not a reliable guide for informing discharge from seclusion.	cli	6	N	N
Hallam KT, Smith DI, Berk M. J Affect Disord. 2009 Jan;112(1-3):212-8. doi: 10.1016/j.jad.2008.03.024. Epub 2008 May 22. Differences between subjective and objective assessments of the utility of Electroconvulsive therapy in patients with bipolar and unipolar depression.	ROM	Use of HoNOS as a clinical measure in ECT for bipolar & unipolar depression	cli	9	N	N
Happell B, Koehn S. Int J Ment Health Nurs. 2010 Jun;19(3):169-76. doi: 10.1111/j.1447-0349.2010.00670.x. From numbers to understanding: the impact of demographic factors on seclusion rates.	ROM	Using HoNOS as one variable in examining seclusion rates	cli	9	N	N
Happell B, Sundram S, Wortans J, Johnstone H, Ryan R, Lakshmana R. Psychiatr Serv. 2009 Nov;60(11):1527-31. doi: 10.1176/appi.ps.60.11.1527. Assessing nurse-initiated care in a mental health crisis assessment and treatment team in Australia.		Evaluation of nurse practitioner	Cli adm	8	N	N
Hasselberg N, Gråwe RW, Johnson S, Ruud T. BMC Health Serv Res. 2011 May 10;11:96. doi: 10.1186/1472-6963-11-96. An implementation study of the crisis resolution team model in Norway: are the crisis resolution teams fulfilling their role?	Res	HoNOS an incidental measure in an evaluation of the CRT model	Cli adm	8	N	N
Hatfield B., Spurrell M., Perry A. Emergency referrals to an acute psychiatric service: Demographic, social and clinical characteristics and comparisons with those receiving continuing services. <i>EmbaseJournal of Mental Health</i> . 9 (3) (pp 305-317), 2000. Date of Publication: 2000	Res	Comparing referrals to existing patients	Cli adm	7	N	N
Hayes RD ¹ , Chang CK, Fernandes AC, Begum A, To D, Broadbent M, Hotopf M, Stewart R. PLoS One. 2012;7(9):e44613. doi: 10.1371/journal.pone.0044613. Epub 2012 Sep 6. Functional status and all-cause mortality in serious mental illness.	ROM	HoNOS factors used as independent variables to predict mortality	cli	9	N	N
Hayes RD, Chang CK, Fernandes A, Begum A, To D, Broadbent M, Hotopf M, Stewart R. J Psychosom Res. 2012 Feb;72(2):114-9. doi:	ROM	Same as above	cli	9	N	N

10.1016/j.jpsychores.2011.09.012. Epub 2011 Nov 20. Associations between symptoms and all-cause mortality in individuals with serious mental illness.						
Hermens MLM, Nugter MA, Mooij N, Hoogendoorn AW, van Son GE, Robbers S, van Bueren MM, Beekman ATF, Theunissen J. Tijdschr Psychiatr. 2017;59(8):456-465. [ROM for the elderly: a prospective cohort study of the outcomes on the HoNOS 65+ in various diagnostic groups]. [Article in Dutch]	cli	justification of old-age psychiatry	adm	7	N	N
Hillier, Bradley; Wright, Lucy; Strydom, Andre; Hassiotis, Angela. Use of the HoNOS-LD in identifying domains of change. <i>PsychINFOThe Psychiatrist</i> . 2014;18(1), 2010, pp. 312-326.	Res	An empirical field trial of the Learning Disabilities version of HoNOS	adm	7	N	N
Honig A, Visser I, Heller H, Kieviet N, Boenink AD. Ned Tijdschr Geneeskd. 2014;158(5):A6520. [A medical-psychiatric unit in a general hospital: effective combined somatic and psychiatric care?]. [Article in Dutch]	Res	Justification of a medical-psychiatric unit in a general hospital	adm	8	N	N
Hooke GR, Page AC. Behav Modif. 2002 Oct;26(5):648-59. Predicting outcomes of group cognitive behavior therapy for patients with affective and neurotic disorders.	Res	As a clinical measure in a follow up study	cli	7	N	N
Howe D, Batchelor S, Coates DD. Early Interv Psychiatry. 2017 Aug;11(4):334-341. doi: 10.1111/eip.12252. Epub 2015 May 12. Young Australians with moderate to severe mental health problems: client data and outcomes at Children and Young People's Mental Health.	ROM	justification of a CAMHS service	adm	8	N	N
Hunter R, Cameron R, Norrie J. Psychiatr Serv. 2009 Feb;60(2):240-5. doi: 10.1176/appi.ps.60.2.240. Using patient-reported outcomes in schizophrenia: the Scottish Schizophrenia Outcomes Study.	ROM	Roms are feasible	cli	9	N	N
Huxley P, Reilly S, Robinshaw E, Mohamad H, Harrison J, Windle B, Butler T. Soc Psychiatry Psychiatr Epidemiol. 2003 Jan;38(1):44-8. Interventions and outcomes of health and social care service provision for people with severe mental illness in England.	ROM	Comparison of socl srvc and mhs's	Adm ali	7	N	N
Jaeger M, Briner D, Kawohl W, Seifritz E, Baumgartner-Nietlisbach. Psychiatry Res. 2015 Dec 15;230(2):413-8. doi: 0.1016/j.psychres.2015.09.029. Epub 2015 Sep 21. Psychosocial functioning of individuals with schizophrenia in community housing facilities and the psychiatric hospital in Zurich.	Cli	Comparison of two populations	cli	7	N	N
Jaeger M, Konrad A, Rueegg S, Rabenschlag F. Nord J Psychiatry. 2015 Apr;69(3):188-95. doi: 10.3109/08039488.2014.959561. Epub 2014 Sep 22. Patients' subjective perspective on recovery orientation on an acute psychiatric unit.	Res	HoNOS used to measure effect of introduction of a recovery project in an IP unit	cli	8	N	N

Jury A, Lai J, Tuason C, Koning A, Smith M, Boyd L, Swanson C, Fergusson D, Gruar A. Int J Ment Health Nurs. 2018 Jul 16. doi: 10.1111/inm.12521. [Epub ahead of print] People who experience seclusion in adult mental health inpatient services: An examination of health of the nation outcome scales scores.	Cli ROM	Correlating individual HoNOS factors with seclusion	Cli	9	N	N
Keet R., Van Veldhuizen J.R., Nugter A. Function assertive community treatment: Treatment of severe mental illness in the Netherlands. <i>EmbaseEuropean Neuropsychopharmacology. Conference: 22 ECNP Congress. Istanbul Turkey. Conference Publication: (var.pagings). 19 (SUPPL. 3) (pp S529), 2009. Date of Publication: 2009.</i>	res	Evaluation of FACT	Cli adm	8	N	N
Kelly C, Sharkey V, Morrison G, Allardyce J, McCreadie RG. Br J Psychiatry. 2000 Oct;177:348-53. Nithsdale Schizophrenia Surveys. 20. Cognitive function in a catchment-area-based population of patients with schizophrenia.	Res	Survey of a community population	Cli adm	7	N	N
Kelly, Peter J; Deane, Frank P. Relationship between therapeutic homework and clinical outcomes for individuals with severe mental illness. <i>PsycINFOAustralian and New Zealand Journal of Psychiatry. Vol.43(10), 2009, pp. 968-975</i>	Res ? ROM ?	deduction from a correlation between homework completion and improvement	cli	9	N	N
Kennedy C, Yellowlees P. J Telemed Telecare. 2003;9(1):12-6. The effectiveness of telepsychiatry measured using the Health of the Nation Outcome Scale and the Mental Health Inventory.	ROM	Part of a comparison of telepsychiatry with face-to-face	Cli adm	8	N	N
Kightley M, Einfeld S, Hancock N. Australas Psychiatry. 2010 Apr;18(2):167-9. doi: 10.3109/10398560903473660. Routine outcome measurement in mental health: feasibility for examining effectiveness of an NGO.	ROM	Inadequate data had been collected to allow assessment of a service	adm	7	N	N
Kilian R., Becker T., Frasc K. Effectiveness and cost-effectiveness of home treatment compared with inpatient care for patients with acute mental disorders in a rural catchment area in Germany. <i>EmbaseNeurology Psychiatry and Brain Research. 22 (2) (pp 81-86), 2016. Date of Publication: 01 Jun 2016</i>	res	Evaluation of HTT	Adm cli	8	N	N
Kisely S, Xiao J, Crowe E, Paydar A, Jian L. Psychiatry Res. 2014 Mar 30;215(3):574-8. doi: 10.1016/j.psychres.2013.12.036. Epub 2013 Dec 30. The effect of community treatment orders on outcome as assessed by the Health of the Nation Outcome Scales.	ROM	CTOs make no difference to outcomes	cli	8	N	N
Koch, Nathalie; Wasserfallen, Jean-Blaise; Camus, Didier; Gebel, Sophia; Bonsack, Charles. Implementing validated routine outcome assessment in psychiatry: A field study. <i>PsycINFOSchweizer Archiv fur Neurologie und Psychiatrie. Vol.164(4), 2013, pp. 134-144</i>	ROM	Swiss field trial	Adm cli	9	N	N

Kodagalli, Anil; Mynors-Wallis, Laurence; Cope, Denise; Ogollah, Reuben; Immins, Tikki. Patient-reported outcome measures v. clinician-measured outcomes in community psychiatric practice. <i>PsycINFOThe Psychiatrist</i> . Vol.36(2), 2012, pp. 61-64	res	Field trial	Adm cli	9	N	N
König HH; Heider D; Rechlin T; Hoffmann P; Birker T; Heinrich S; Brettschneider C; Hierholzer C; Riedel-Heller SG; Roick C. [How does the Regional Psychiatry Budget (RPB) work in an area with initially low capacity of psychiatric hospital beds?]. [German] <Wie wirkt das Regionale Psychiatriebudget (RPB) in einer Region mit initial niedriger Bettenmessziffer?> <i>Ovid MEDLINE(R)Psychiatrische Praxis</i> . 40(8):430-8, 2013 Nov	res	HoNOS a clinical measure in a study of a new funding model	adm	8	N	N
König, Hans-Helmut; Heinrich, Sven; Heider, Dirk; Deister, Arno; Zeichner, Dirk; Birker, Thomas; Hierholzer, Cornelia; Angermeyer, Matthias C; Roick, Christiane. The regional psychiatry budget (RPB): A model for a new payment system of hospital based mental health care services? Analysis of costs and effectiveness of the RPB at 3.5 years follow-up. [German]. Das regionale psychiatriebudget (RPB): Ein modell für das neue pauschalierende entgeltsystem psychiatrischer krankenhausleistungen? Analyse der kosten und effekte des RPB nach 3,5 jahren laufzeit. <i>PsycINFOPsychiatrische Praxis</i> . Vol.37(1), 2010, pp. 34-42	res	As above	adm	8	N	N
König HH, Heinrich S, Heider D, Deister A, Zeichner D, Birker T, Hierholzer C, Angermeyer MC, Roick C. Psychiatr Prax. 2010 Jan;37(1):34-42. doi: 10.1055/s-0029-1223418. Epub 2010 Jan 13. [The regional psychiatry budget (RPB): a model for a new payment system of hospital based mental health care services]. [Article in German]		Used as a measure in the assessment of a new method of financing services	adm	8	N	N
Kortrijk HE, Mulder CL, Drukker M, Wiersma D, Duivenvoorden HJ. Aust N Z J Psychiatry. 2012 Mar;46(3):240-8. doi: 10.1177/0004867411433214. Epub 2012 Jan 5. Duration of assertive community treatment and the interpretation of routine outcome data.	ROM	ACT outcomes vary with treatment duration	adm	7	N	N
Kortrijk HE, Mulder CL, Roosenschoon BJ, Wiersma D. Community Ment Health J. 2010 Aug;46(4):330-6. doi: 10.1007/s10597-009-9257-9. Epub 2009 Oct 22. Treatment outcome in patients receiving assertive community treatment.	ROM	Fairly similar conclusions and purpose to the previous study	adm	7	N	N
Kortrijk HE, Staring AB, van Baars AW, Mulder CL. Soc Psychiatry Psychiatr Epidemiol. 2010 Feb;45(2):245-52. doi: 10.1007/s00127-009-0061-1. Epub 2009 May 2. Involuntary admission may support treatment outcome and motivation in patients receiving assertive community treatment.	ROM	A measure in assessing involuntary treatment during the course of ACT	cli	8	N	N

Kotynia-English R ¹ , McGowan H, Almeida OP. Int Psychogeriatr. 2005 Sep;17(3):475-85. A randomized trial of early psychiatric intervention in residential care: impact on health outcomes.	ROM	Screening and early referral did not improve outcomes	Cli adm	8	N	N
Lai J, Jury A, Long J, Fergusson D, Smith M, Baxendine S, Gruar A Variation in seclusion rates across New Zealand's specialist mental health services: Are sociodemographic and clinical factors influencing this? Int J Ment Health Nurs. 2018 Aug 18	Cli ROM	As a controlling variable in study	Adm & cli	9	N	N
Lakeman R, Bradbury J. J Psychiatr Ment Health Nurs. 2014 May;21(4):327-35. doi: 10.1111/jpm.12089. Epub 2013 May 22. Mental health nurses in primary care: quantitative outcomes of the Mental Health Nurse Incentive Program.	res	Evaluation of mental health nurses in primary care	Adm cli	8	N	N
Lanfredi M., De Girolamo G., Candini V., Buizza C., Ferrari C., Boero M.E., Giobbio G.M., Goldschmidt N., Greppo S., Iozzino L., Maggi P., Melegari A.L., Pasqualetti P., Rossi G. Predictors of quality of life in a sample of inpatients with schizophrenia in 23 residential facilities in northern Italy. <i>EmbaseEuropean Psychiatry. Conference: 21st European Congress of Psychiatry, EPA 2013. Nice France. Conference Publication: (var.pagings). 28 (SUPPL. 1) (no pagination), 2013. Date of Publication: 2013.</i>	Res	Used as a clinical measure in a one year longitudinal survey	Adm cli	7	N	N
Laugharne R, Eaves S, Mascas A, Psatha K, Dinnis G, Trower J, Shankar R. BJPsych Open. 2018 May;4(3):91-94. doi: 10.1192/bjo.2018.3. Desperately seeking outcomes: quantifying the effectiveness of community mental healthcare using Health of the Nation Outcome Scales.	Adm	measure of which patients do better in CMHTs and which do not	cli	8	N	N
Long, Clive G; Dickens, Geoff; Sugarman, Philip; Craig, Lorraine; Mochty, Ursula; Hollin, Clive. Tracking risk profiles and outcome in a medium secure service for women: Use of the HoNOS-Secure. [References]. <i>PsycINFOThe International Journal of Forensic Mental Health. Vol.9(3), 2010, pp. 215-225.</i>	ROM	Review of 5 years of outcomes measures – authors assert that this enabled transfer to less secure settings (but no attempt to compare with decision-making without rom)	cli	6	N	N
Lopez-Morinigo JD, Fernandes AC, Chang CK, Hayes RD, Broadbent M, Stewart R, David AS, Dutta. RBMC Psychiatry. 2014 Aug 1;14:213. doi: 10.1186/s12888-014-0213-z. Suicide completion in secondary mental healthcare: a comparison study between schizophrenia spectrum disorders and all other diagnoses.	ROM	Comparison of suicides of patients with and without schizophrenia	cli	7	N	N
Lora A, Bezzi R, Erlicher A. Community Ment Health J. 2007 Aug;43(4):341-57. Epub 2007 Jan 26. Estimating the prevalence of severe mental illness in mental health services in Lombardy (Italy).	Res ? ROM ?	What the title says	Cli adm	9	N	N

Lorant V, Grard A, Van Audenhove C, Helmer E, Vanderhaegen J, Nicaise P. BMC Health Serv Res. 2016 May 17;16:187. doi: 10.1186/s12913-016-1434-2. Assessment of the priority target group of mental health service networks within a nation-wide reform of adult psychiatry in Belgium.	Res	HoNOS (& other measures) used to compare patients accepted or not accepted by services	adm	7	N	N
Luo W, Harvey R, Tran T, Phung D, Venkatesh S, Connor JP. BMJ Open. 2016 Apr 27;6(4):e010732. doi: 10.1136/bmjopen-2015-010732. Consistency of the Health of the Nation Outcome Scales (HoNOS) at inpatient-to-community transition.	ROM	Transfer from IP to community setting magically increased the HoNOS score	Cli adm	9	N	N
Macpherson R, Calciu C, Foy C, Humby K, Lozynskyj D, Garton C, Steer H, Elliott H. BJPsych Bull. 2017 Dec;41(6):330-336. doi: 10.1192/pb.bp.116.055137. A service evaluation of outcomes in two in-patient recovery units.	Cli	Evaluation of en-patient recovery units	cli	7	N	N
Macpherson, Rob; Peddu, Pradeep; David, Chris. The Vron - An early intervention in-patient recovery unit. <i>PsycINFOIrish Journal of Psychological Medicine. Vol.29(2), 2012, pp. 113-116</i>	ROM	A clinical measure in a new model of IP care	Cli adm	8	N	N
Macpherson R, Pesola F, Leamy M, Bird V, Le Boutillier C, Williams J, Slade M. Schizophr Res. 2016 Aug;175(1-3):142-147. doi: 10.1016/j.schres.2015.10.031. Epub 2015 Oct 31. The relationship between clinical and recovery dimensions of outcome in mental health.	ROM	Use of HoNOS to help differentiate between clinical and personal recovery	cli	9	N	N
Macpherson, Rob; Thyarappa, Praveen; Riley, Genevieve; Steer, Hannah; Blackburn, Mike; Foy, Chris. Evaluation of three assertive outreach teams. <i>PsycINFOThe Psychiatrist. Vol.37(7), 2013, pp. 228-231</i>	ROM	What the title says	Cli adm	8	N	N
Maddison P, Golay P, Muncer SJ. Eur Psychiatry. 2018 Feb;48:1-5. doi: 10.1016/j.eurpsy.2017.10.006. Epub 2018 Jan 10. Latent class analysis of the health of the nation outcome scales: A comparison of Swiss and English profiles and exploration of their predictive utility.	Cli	Study of the predictive utility of HoNOS in UK and Swiss settings	res	9	N	N
Mapelli V., Bezzi R., Erlicher A., Lora A., Miragoli P. Cost-effectiveness of treatment of schizophrenia with typical and atypical antipsychotics. <Costi ed efficacia della cura della schizofrenia con antipsicotici tipici e atipici.> <i>EmbasePharmacoEconomics - Italian Research Articles. 6 (1) (pp 15-30), 2004. Date of Publication: 2004</i>	Res	cost comparison of different antipsychotics.	cli	7	N	
McColl H, Johnson S. Soc Psychiatry Psychiatr Epidemiol. 2006 Oct;41(10):789-95. Epub 2006 Aug 9. Characteristics and needs of asylum seekers and refugees in contact with London community mental health teams: a descriptive investigation.	ROM	Descriptive survey of asylum seekers	adm	7	N	N
McCrossan P, Ryan A, Connellan M, Power P. Ir J Psychol Med. 2017 Mar;34(1):39-44. doi: 10.1017/ipm.2016.12. The impact of a specialized	res	Evaluation of a young adults programme (transition from CAMHS to adult)	adm	8	N	N

inpatient and day patient group programme on clinical outcome in older adolescents and young adults with mental illness.						
McKay R, Coombs T, Duerden D. Australas Psychiatry. 2014 Feb;22(1):13-8. doi: 10.1177/1039856213511673. Epub 2013 Nov 27. The art and science of using routine outcome measurement in mental health benchmarking.	ROM	described bi-annual benchmarking forums in New South Wales, Australia, in which certain aspects of outcome measures were provided for comparison of services. This has not been rolled out at a national level. The impacts of these forums on clinical practice were not assessed. described bi-annual benchmarking forums in New South Wales, Australia, in which certain aspects of outcome measures were provided for comparison of services. This has not been rolled out at a national level. The impacts of these forums on clinical practice were not assessed.	Cli adm	9	N	N
McLeod HJ, Jaques S, Deane FP. Psychiatr Rehabil J. 2009 Spring;32(4):269-75. doi: 10.2975/32.4.2009.269.275. Base rates of physical activity in Australians with schizophrenia.	ROM	x-sectional survey of patients	cli	9	N	N
Meagher, David; O'Brien, Sinead; Pulella, Ananth; Oshun, Anthony; Brosnan, Pat. Multidisciplinary activities in a community mental health service: Relationship to Health of the Nation Outcome Scale scores and diagnosis. <i>PsycINFO Psychiatric Bulletin</i> . Vol.33(5), 2009, pp. 172-175.	res	HoNOS a variable in description of patients of a CMHS	Cli adm	7	N	N
Mellsop G, Smith B. Aust N Z J Psychiatry. 2007 Oct;41(10):850-3. Reflections on masculinity, culture and the diagnosis of depression.	ROM	Used as clinical measure in study of a depressed patients	cli	9	N	N
Meneghelli, Anna; Breda, Alessandro; Caprin, Claudia; Patelli, Giovanni; Cocchi, Angelo. Early intervention in psychosis: Assessment modalities. [Italian]. L'intervento precoce nelle psicosi: Strategie di valutazione. <i>PsycINFO Psicoterapia Cognitiva e Comportamentale</i> . Vol.9(1), 2003, pp. 17-30	Res	Part of an assessment battery – patient characteristics reported	cli	7	N	N
Meneghelli A., Cocchi A. Programma2000: Concept and practical experiences. <i>Embase European Archives of Psychiatry and Clinical Neuroscience. Conference: 3rd European Conference on Schizophrenia Research: Facts and Visions. Berlin Germany. Conference Publication: (var.pagings). 261 (SUPPL. 1) (pp S37), 2011. Date of Publication: September 2011.</i>	Res	Evaluation of eip programme	Cli adm	8	N	N
Meneghelli A., Cocchi A. Treatment, treatment UHR, cannabis use Programma 2000: A leading-edge programme on early psychosis. <i>Embase Early Intervention in Psychiatry. Conference: 7th International</i>	Res	ditto	Cli adm	8	N	N

Conference on Early Psychosis - Early Psychoses: A Lifetime Perspective. Amsterdam Netherlands. Conference Publication: (var.pagings). 4 (SUPPL. 1) (pp 6), 2010. Date of Publication: November 2010						
Meneghelli A., Patelli G., Cocchi A., Preti A. Investigating expressed emotion in families of FEP and UHR schizophrenia patients: PROGRAMMA2000 findings. <i>EmbaseEarly Intervention in Psychiatry. Conference: 8th International Conference on Early Psychosis: From Neurobiology to Public Policy. San Francisco, CA United States. Conference Publication: (var.pagings). 6 (SUPPL.1) (pp 91), 2012. Date of Publication: October 2012.</i>	Res	Follow up study of EIP	cli	7	N	N
Mirabella, Fiorino; Rosicarelli, Maria Laura; Broccoli, Giorgio; Leoncini, Luciano; Marchesini, Nadia; Morosini, Pierluigi. Effectiveness assessment of a structured cognitive-behavioural group therapy on psychiatric inpatients in day centres. [Italian]. Valutazione dell'efficacia di un intervento strutturato cognitivo-comportamentale di gruppo su pazienti psichiatrici ospiti di Centri Diurni. <i>PsycINFOPsicoterapia Cognitiva e Comportamentale. Vol.15(3), 2009, pp. 329-352</i>	Res	Evaluation of CBT	Cli adm	8	N	N
Monzani E, Erlicher A, Lora A, Lovaglio P, Vittadini G. Int J Ment Health Syst. 2008 Jul 5;2(1):10. doi: 10.1186/1752-4458-2-10. Does community care work? A model to evaluate the effectiveness of mental health services.	ROM	comparison of different services	adm	8	N	N
Moran V, Jacobs R. Eur J Health Econ. 2018 Jun;19(5):709-718. doi: 10.1007/s10198-017-0915-5. Epub 2017 Jun 24. Investigating the relationship between costs and outcomes for English mental health providers: a bi-variate multi-level regression analysis.	ROM	Evaluation of cost containment in UK	Cli adm	7	N	N
Moran V, Jacobs R. Soc Sci Med. 2015 Sep;140:127-35. doi: 10.1016/j.socscimed.2015.07.009. Epub 2015 Jul 13. Comparing the performance of English mental health providers in achieving patient outcomes.	ROM	Comparing services by outcome Different services have different outcomes and will therefore receive different levels of payment.	adm	9	N	N
Mozley CG, Huxley P, Sutcliffe C, Bagley H, Burns A, Challis D, Cordingley L. Int J Geriatr Psychiatry. 1999 Sep;14(9):776-83. 'Not knowing where I am doesn't mean I don't know what I like': cognitive impairment and quality of life responses in elderly people.	Res	HoNOS part of a battery to tests to assess how well cognitively impaired patients can articulate their quality of life	cli	9	N	N
Müller M, Vandeleur C, Weniger G, Prinz S, Vetter S, Egger ST. Psychiatry Res. 2016 May 30;239:20-7. doi: 10.1016/j.psychres.2016.02.060. Epub 2016 Mar 3. The performance of the Health of the Nation Outcome Scales as measures of clinical severity.	Rom (pres umab ly)	Assessment of HoNOS as a measure of clinical severity	Cli adm	9	N	N

Mueller-Stierlin AS, Helmbrecht MJ, Herder K, Prinz S, Rosenfeld N, Walendzik J, Holzmann M, Dinc U, Schützwohl M, Becker T, Kilian R. BMC Psychiatry. 2017 Aug 1;17(1):283. doi: 10.1186/s12888-017-1441-9. Does one size really fit all? The effectiveness of a non-diagnosis-specific integrated mental health care program in Germany in a prospective, parallel-group controlled multi-centre trial.	res	Evaluation of treatment delivery programme	adm	8	N	N
Mulligan O., Helmi L., Adamis D., O'Mahony E., McCarthy G., MacDonald A. Implementation of Health of the Nation Outcome Scale (HoNOS) in outpatient clinic, sligo mental health services: Feasibility and agreement with global assessment of functioning scale. <i>EmbaseEuropean Psychiatry. Conference: 24th European Congress of Psychiatry, EPA 2016. Madrid Spain. Conference Publication: (var.pagings). 33 (SUPPL.) (pp S483), 2016. Date of Publication: March 2016</i>	ROM	HoNOS showed that patients improved.	Adm+ cli	9	N	N
Mujic F, Cairns R, Mak V, Squire C, Wells A, Al-Harrasi A, Prince M. BJPsych Bull. 2018 Feb;42(1):30-36. doi: 10.1192/bjb.2017.9. Liaison psychiatry for older adults in the general hospital: service activity, development and outcomes.	Cli	justification of liaison psychiatry in older adults	adm	7	N	N
Mulder CL, Jochems E, Kortrijk HE. Soc Psychiatry Psychiatr Epidemiol. 2014 Apr;49(4):541-8. doi: 10.1007/s00127-013-0779-7. Epub 2013 Oct 18. The motivation paradox: higher psychosocial problem levels in severely mentally ill patients are associated with less motivation for treatment.	Res ROM	Motivation for treatment in patients with high HoNOS scores	cli	9	N	N
Mulder C.L., Kortrijk H. Functional and symptomatic remission and its association with quality of life. <i>EmbaseEarly Intervention in Psychiatry. Conference: 7th International Conference on Early Psychosis - Early Psychoses: A Lifetime Perspective. Amsterdam Netherlands. Conference Publication: (var.pagings). 4 (SUPPL. 1) (pp 19), 2010. Date of Publication: November 201</i>	Res	Clinical measure looking at the title	cli	7	N	N
Muncer S, Bass M, Dawkin M. Australas Psychiatry. 2016 Oct;24(5):459-61. doi: 10.1177/1039856216647254. Epub 2016 Apr 28. Mokken analysis of the Health of the Nation Outcome Scales in acute inpatient and community samples.	Res	Comparison of HoNOS with Mokken analysis in a cohort of patients	Cli adm	9	N	N
Muncer S, Speak B. Psychiatry Res. 2017 Jan;247:12-14. doi: 10.1016/j.psychres.2016.11.010. Epub 2016 Nov 12. Confirmatory factor analysis of a two scale model of the Health of the Nation Outcome Scales (HoNOS) across diagnostic categories.	ROM	How well does HoNOS fit with the cluster groupings?	adm	7	N	N
Munz I, Ott M, Jahn H, Rauscher A, Jäger M, Kilian R, Frasch K. Psychiatr Prax. 2011 Apr;38(3):123-8. doi: 10.1055/s-0030-1248591. Epub 2010 Sep	Res	Evaluation of home treatment	Cli adm	8	N	N

16. [Comparison of "home treatment" with traditional inpatient treatment in a mental hospital in rural southern Germany]. [Article in German]						
Murphy D, Hodgman G, Carson C, Spencer-Harper L, Hinton M, Wessely S, Busuttill W. BMJ Open. 2015 Mar 20;5(3):e007051. doi: 10.1136/bmjopen-2014-007051. Mental health and functional impairment outcomes following a 6-week intensive treatment programme for UK military veterans with post-traumatic stress disorder (PTSD): a naturalistic study to explore dropout and health outcomes at follow-up.	ROM	justification of an intensive 6-week PTSD programme with veterans	adm	8	N	N
Cognitive behaviour therapy for schizophrenia in patients with mild to moderate substance misuse problems. [References]. Naeem, Farooq; Kingdon, David; Turkington, Douglas. <i>PsycINFO Cognitive Behaviour Therapy</i> . Vol.34(4), 2005, pp. 207-215	res	Evaluation of cbt	Adm cli	8	N	N
Naeem, Farooq; Bhatti, Faiz; Pickering, Ruth; Kingdon, David. A controlled trial of the effectiveness of drug treatment & testing orders (DTTO) with standard care. <i>PsycINFO Journal of Substance Use</i> . Vol.12(4), 2007, pp. 253-265	Res ? ROM	Evaluation of outcome of such orders	Cli adm	8	N	N
Naeem, Farooq; Kingdon, David; Turkington, Douglas. Predictors of response to cognitive behaviour therapy in the treatment of schizophrenia: A comparison of brief and standard interventions. <i>PsycINFO Cognitive Therapy and Research</i> . Vol.32(5), 2008, pp. 651-656	Res	A clinical measure in assessing CBT	Cli adm	8	N	N
Nayar B., Patel M., Grant W. Joint Perinatal Mental Health Clinic (PMH) in North West London hospital-a success story of collaborative working. <i>EmbaseBJOG: An International Journal of Obstetrics and Gynaecology</i> . Conference: RCOG World Congress 2016. United Kingdom. 123 (Supplement 2) (pp 141-142), 2016. Date of Publication: June 2016	Res	Evaluation of embedding a psychiatrist in a perinatal service	Cli adm	8	N	N
O'Connor N., Lambert T., Mulley R., Grasso B. HoNOS vs MIR-S-CGI: data from an Acute Recovery Unit. <i>Embase Australian and New Zealand Journal of Psychiatry</i> . Conference: Royal Australian and New Zealand College of Psychiatrists, RANZCP Annual Congress 2011. Darwin, NT Australia. Conference Publication: (var.pagings). 45 (O1) (pp A47), 2011. Date of Publication: May 2011	ROM	Used as a severity measure in assessing patients' response to treatment	Cli adm	8	N	N
Meagher D, O'Brien S, McFarland J, Kealy B, Pullela A, Saunders J, Cullen W., Ir J Med Sci. 2012 Sep;181(3):301-8. doi: 10.1007/s11845-011-0795-0. Epub 2012 Jan 5. A randomized-controlled trial of intensive case management emphasizing the recovery model among patients with severe and enduring mental illness.	Res ROM ?	Evaluation of the introduction of recovery principles	cli	8	N	N
Øiesvold T, Bakkejord T, Hansen V, Nivison M, Sørgaard KW.	res	Casual baseline measure in a study	cli	9	N	N

Soc Psychiatry Psychiatr Epidemiol. 2012 Mar;47(3):419-25. doi: 10.1007/s00127-011-0343-2. Epub 2011 Feb 2. Suicidality related to first-time admissions to psychiatric hospital.						
Okruashvili M., Roinishvili M., Brand A., Herzog M.H., Chkonja E. Predictors of better social functioning for patients with first episode psychoses. <i>EmbaseEuropean Psychiatry. Conference: 23rd European Congress of Psychiatry, EPA 2015. Vienna Austria. Conference Publication: (var.pagings). 30 (SUPPL. 1) (pp 269), 2015. Date of Publication: 31 Mar 2015.</i>	Res	FEP study – clinical tool	cli	7	N	N
Olso, Turid Moller; Gudde, Camilla Buch; Wullum, Elin; Linaker, Olav M. Gender-specific correlates of substance use in patients with serious mental illness. <i>PsycINFOMental Health and Substance Use. Vol.5(1), 2012, pp. 31-41</i>	Res	Same as the next one	cli	9	N	N
Olso, Turid Moller; Gudde, Camilla Buch; Wullum, Elin; Linaker, Olav M. Associations between symptoms, functioning and self-reported motivations for alcohol use and alcohol problems in patients with serious mental illness. <i>PsycINFOMental Health and Substance Use. Vol.6(3), 2013, pp. 237-249.</i>	Res	HoNOS used as a measure in examination of alcohol use	cli	9	N	N
Østergaard SD, Bille J, Søltoft-Jensen H, Lauge N, Bech P.J Affect Disord. 2012 Sep;140(1):48-56. doi: 10.1016/j.jad.2012.01.039. Epub 2012 Mar 3. The validity of the severity-psychosis hypothesis in depression.	Res ROM	Using HoNOS to tease out differences between psychotic and non-psychotic depression	cli	9	N	N
Page AC, Hooke GR. J Affect Disord. 2004 Sep;81(3):275-8. Failure to replicate the effects of gender and season on the length of hospitalisation in unipolar depressives.	ROM	Use of one scale of HoNOS as a clinical measure	cli	9	N	N
James, Michael; Kehoe, Robert. Using the Health of the Nation Outcome Scales in clinical practice. <i>PsycINFOPsychiatric Bulletin. Vol.23(9), 1999, pp. 536-538</i>	Res	a description of how they set up the use of HoNOS – no attempt to determine whether it makes any difference	Cli adm	9	N	N
Palic S, Kappel ML, Nielsen MS, Carlsson J, Bech P. BMC Psychiatry. 2014 Dec 18;14:330. doi: 10.1186/s12888-014-0330-8. Comparison of psychiatric disability on the health of nation outcome scales (HoNOS) in resettled traumatized refugee outpatients and Danish inpatients.	ROM	Using HoNOS to compare different patient populations	cli	9	N	N
Parabiaghi A, Kortrijk HE, Mulder CL. Soc Psychiatry Psychiatr Epidemiol. 2014 Feb;49(2):291-305. doi: 10.1007/s00127-013-0750-7. Epub 2013 Aug 6. Defining multiple criteria for meaningful outcome in routine outcome measurement using the Health of the Nation Outcome Scales.	ROM	Value of HoNOS as a clinical measure is tested in clinical samples in 2 different countries. Different guidelines for interpreting HoNOS findings gave different results	cli	9	N	N
Parabiaghi A, Rapisarda F, D'Avanzo B, Erlicher A, Lora A, Barbato A. Aust N Z J Psychiatry. 2011 Jul;45(7):558-68. doi: 0.3109/00048674.2011.580450. Epub 2011 May 11. Measuring clinical	Res	Promoting ROMS but not providing any evidence of their value	Cli adm	9	N	N

change in routine mental health care: differences between first time and longer term service users.						
Patel R, Oduola S, Callard F, Wykes T, Broadbent M, Stewart R, Craig TK, McGuire P. BMJ Open. 2017 Mar 9;7(3):e013113. doi: 10.1136/bmjopen-2016-013113. What proportion of patients with psychosis is willing to take part in research? A mental health electronic case register analysis.	ROM	Correlation of data with willingness to participate in research	res	9	N	N
Pec O, Bob P, Pec J, Hrubcova A. Psychol Psychother. 2018 Jun;91(2):157-168. doi: 10.1111/papt.12153. Epub 2017 Sep 13. Psychodynamic day treatment programme for patients with schizophrenia spectrum disorders: Dynamics and predictors of therapeutic change.	Cli	Evaluation of psychodynamic psychotherapy in schizophrenia	cli	8	N	N
Pec O, Bob P, Raboch J. Neuropsychiatr Dis Treat. 2014 Mar 17;10:487-91. doi: 10.2147/NDT.S57627. eCollection 2014. Dissociation in schizophrenia and borderline personality disorder.	Res	Comparison of dissociative symptoms in schizophrenia and BPD	cli	9	N	N
Pec O, Bob P, Raboch J. PLoS One. 2014 Mar 6;9(3):e91228. doi: 10.1371/journal.pone.0091228. eCollection 2014. Splitting in schizophrenia and borderline personality disorder.	Res	Comparison of splitting in schizophrenia and BPD	cli	9	N	N
Penno SJ, Hamilton B, Petrakis M. Arch Psychiatr Nurs. 2017 Dec;31(6):553-560. doi: 10.1016/j.apnu.2017.07.003. Epub 2017 Jul 17. Early Intervention in Psychosis: Health of the Nation Outcome Scales (HoNOS) Outcomes From a Five-Year Prospective Study.	Cli	evaluation of EIP	adm	8	N	N
Percudani M, Gerzeli S, Massagrani R, Jommi C, Fattore G, Cerati G, Contini A. Epidemiol Psichiatri Soc. 2003 Jul-Sep;12(3):175-86. [Costs and outcome of care in subjects with severe mental disorders]. [Article in Italian]	Res	Correlating severity measures at admission with overall treatment costs	Adm cli	7	N	N
Pillay, Selena M; Oliver, Brid; Butler, Louise; Kennedy, Harry G. Risk stratification and the care pathway. <i>PsycINFOIrish Journal of Psychological Medicine. Vol.25(4), 2008, pp. 123-127</i>	res	HoNOS used as a clinical measure in the assessment of a stratified model of forensic care	Cli adm	8	N	N
Pirkis J, Burgess P, Jolley D. Soc Psychiatry Psychiatr Epidemiol. 1999 Dec;34(12):634-44. Suicide attempts by psychiatric patients in acute inpatient, long-stay inpatient and community care.	Res	Correlating with suicide attempt incidence	Cli adm	9	N	N
Placentino A, Lucchi F, Scarsato G, Fazzari G; Gruppo REX.IT. Riv Psichiatri. 2017 Nov-Dec;52(6):247-254. doi: 10.1708/2846.28728. [Mental Health Recovery Star: features and validation study of the Italian version]. [Article in Italian]	Res	HoNOS used as comparator for a new evaluation instrument	res	9	N	N
Pompili M, Innamorati M, Gonda X, Serafini G, Erbutto D, Ricci F, Fountoulakis KN, Lester D, Vazquez G, Rihmer Z, Amore M, Girardi P. Hum Psychopharmacol. 2014 Nov;29(6):578-88. doi: 10.1002/hup.2445. Epub 2014 Nov 4. Pharmacotherapy in bipolar disorders during	Res	HoNOS score used as the dependent variable in study	cli	9	N	N

hospitalization and at discharge predicts clinical and psychosocial functioning at follow-up.						
Pompili M, Innamorati M, Gonda X, Serafini G, Sarno S, Erbuto D, Palermo M, Elena Seretti M, Stefani H, Lester D, Perugi G, Akiskal H, Siracusano A, Rihmer Z, Tatarelli R, Amore M, Girardi P. J Affect Disord. 2013 Sep 5;150(2):216-22. doi: 10.1016/j.jad.2013.03.026. Epub 2013 May 17. Affective temperaments and hopelessness as predictors of health and social functioning in mood disorder patients: a prospective follow-up study.	Res	A dependent variable in a study of hopelessness and temperament in bipolar disorder	cli	9	N	N
Prabhu R, Oakley Browne M. Australas Psychiatry. 2008 Jun;16(3):195-9. doi: 10.1080/10398560701784813. The use of the Health of the Nation Outcome Scale in an outreach rehabilitation program.	ROM	HoNOS demonstrates improvement in a rehab programme	adm	8	N	n
Prowse L, Coombs T. Aust Health Rev. 2009 Feb;33(1):13-8. The use of the Health of the Nation Outcome Scales (HoNOS) to inform discharge and transfer decisions in community mental health services.	ROM	commented on in paper	cli	6	N	N
Rees, Anne; Richards, Ann; Shapiro, David A. Utility of the HoNOS in measuring change in a Community Mental Health Care population. <i>PsycINFO Journal of Mental Health. Vol.13(3), 2004, pp. 295-30</i>	ROM	Survey of rom scores in community patients	cli	9	N	N
Rensen YCM, Egger JIM, Westhoff J, Walvoort SJW, Kessels RPC. Int Psychogeriatr. 2018 Jun 20;1-9. doi: 10.1017/S1041610218000492. [Epub ahead of print] The effect of errorless learning on psychotic and affective symptoms, as well as aggression and apathy in patients with Korsakoff's syndrome in long-term care facilities.	Res	A measure of improvement in a trial of errorless learning in Korsakoff's	cli	8	N	N
Restek-Petrović B, Majdančić A, Molnar S, Grah M, Ivezić E, Filipčić I, Bogović A, Grošić V, Mayer N, Kezić S, Pavlović I. Psychiatr Danub. 2017 Jun;29(2):162-170. Early intervention programme for patients with psychotic disorders in "Sveti Ivan" Psychiatric hospital (RIPEPP) - sociodemographic and baseline characteristics of the participants.	Cli	Descriptive study of patients admitted to a particular facility	cli	9	N	N
Ribeiro RB, Tully J, Fotiadou M. Int J Law Psychiatry. 2015 Mar-Apr;39:83-9. doi: 10.1016/j.ijlp.2015.01.025. Epub 2015 Mar 5. Clinical characteristics and outcomes on discharge of women admitted to a Medium Secure Unit over a 4-year period.	ROM	Descriptive study of medium-secure femals patients and naive description of outcomes	Cli Adm	7	N	N
Riguzzi M., Wieser S., Dettling M., Pletscher M. Design of a prospective payment system for inpatient psychiatric care in Switzerland. <i>EmbaseJournal of Mental Health Policy and Economics. Conference: 12th Workshop on Costs and Assessment in Psychiatry. Venice Italy. Conference Publication: (var.pagings). 18 (SUPPL. 1) (pp S33), 2015. Date of Publication: 2015</i>	Res	Evaluation of a new payment system	Adm	8	N	N

Rodenburg H, Bos V, O'Malley C, McGeorge P, Love T, Dowell A. N Z Med J. 2004 Sep 24;117(1202):U1077. General Practice care of enduring mental health problems: an evaluation of the Wellington Mental Health Liaison Service.	Res	Follow up Study of transfer to GP care	cli	8	N	N
Rossi F, Blaco R, Castelli C, Civiotti G, Cocchi A, Contini A, Erlicher A, Lanzara D, Lora A, Marcandalli S, Morosini P, Pisapia D. Epidemiol Psichiatr Soc. 1999 Jul-Sep;8(3):198-208. [Cost of psychiatric patients by disability type]. [Article in Italian]	Res	Cost of care study	adm	8	N	N
Rushforth D., Brooker C., Winstanley J., Reppe D. The use of HoNOS in the evaluation of partial hospitalization in mental health day care. <i>Ovid EmcareClinical Effectiveness in Nursing</i> . 4 (3) (pp 121-127), 2000. Date of Publication: September 200	Res	Evaluation of partial hospitalisation	Cli adm	8	N	N
Ruud T, Aarre TF, Boeskov B, le Husevåg PS, Klepp R, Kristiansen SA, Sandvik J. Int J Ment Health Syst. 2016 Apr 12;10:33. doi: 10.1186/s13033-016-0064-8. eCollection 2016. Satisfaction with primary care and mental health care among individuals with severe mental illness in a rural area: a seven-year follow-up study of a clinical cohort.	Res	Almost incidental used of HoNOS in a satisfaction survey	Cli adm	9	N	N
Ryan, Sean. Strengths-based approach in psychotherapy with intellectually disabled individuals. <i>PsycINFO Dissertation Abstracts International: Section B: The Sciences and Engineering</i> . Vol.77(1-B(E)),2016, pp. No Pagination Specified.	Res	Clinical measure in an outcomes study	cli	9	N	N
Samanes R.O., Tejedor A.G., Muro C. Social integration of institutionalized elderly. <Integración social del anciano institucionalizado.> <i>EmbaseGerokomos</i> . 23 (4) (pp 172-176), 2012. Date of Publication: December 2012.	Res	Part of the assessment of a specific population	Adm cli	7	N	N
Sancassiani F, Lorrai S, Cossu G, Cocco A, Trincas G, Floris F, Mellino G, Machado S, Nardi AE, Fabrici EP, Preti A, Carta MG. Clin Pract Epidemiol Ment Health. 2017 Nov 24;13:220-232. doi: 10.2174/1745017901713010220. eCollection 2017. The Effects of "VelaMente?!" Project on Social Functioning of People With Severe Psychosocial Disabilities.	Res	Evaluation of sailing in a crew as therapy	cli	8	N	N
Schneider, Justine; Brandon, Toby; Wooff, David; Carpenter, John; Paxton, Roger. Assertive outreach: Policy and reality. <i>PsycINFO Psychiatric Bulletin</i> . Vol.30(3), 2006, pp. 89-94.	ROM	Evaluation of assertive outreach teams	Adm cli	8	N	N
Schulte P., Veerman S., Smith J., De Haan L. Memantine augmentation in clozapine refractory schizophrenia: A randomized, double-blind, placebo-controlled crossover study. <i>EmbaseEuropean Archives of Psychiatry and Clinical Neuroscience</i> . Conference: 5th European Conference on	Res	Clinical measure in drug trial	cli	8	N	N

<i>Schizophrenia Research: Bridging Gaps Improving Outcomes. Berlin Germany. Conference Publication: (var.pagings). 265 (1 SUPPL. 1) (pp S66), 2015. Date of Publication: September 2015</i>						
Segal SP, Hayes SL, Rimes L. Psychiatr Serv. 2017 Dec 1;68(12):1247-1254. doi: 10.1176/appi.ps.201600161. Epub 2017 Aug 1. The Utility of Outpatient Commitment: I. A Need for Treatment and a Least Restrictive Alternative to Psychiatric Hospitalization.	ROM	Comparison of needs of CTO with non-CTO patients	cli	8	N	N
Sender-Galloway, Simeon; Clark, Tom. Problematic substance use among forensic psychiatric community patients. <i>PsycINFO Journal of Forensic Practice. Vol.15(2), 2013, pp. 119-129</i>	Res ? ROM ?	Used as a clinical measure in a study of substance use	Cli adm	7	N	N
Sharma VK, Wilkinson G, Fear S. Br J Psychiatry. 1999 May;174:395-8. Health of the Nation Outcome Scales: a case study in general psychiatry.	Cli	Field trial	res	9	N	N
Shed J., Priest P. The Health of the Nation Outcome Scale Payment by Results (HoNOS PbR): Clinician and service user perspectives on the tool. <i>Ovid Emcare Clinical Psychology Forum. (255) (pp 32-36), 2014. Date of Publication: March 2014</i>	unkn own	"a small scale survey"	unkno wn	9	N	N
Siskind D, Harris M, Kisely S, Siskind V, Brogan J, Pirkis J, Crompton D, Whiteford H. Community Ment Health J. 2014 Jul;50(5):538-47. doi: 10.1007/s10597-013-9654-y. Epub 2013 Oct 23. A retrospective quasi-experimental study of a transitional housing program for patients with severe and persistent mental illness.	ROM	Evaluation of a transitional housing programme	Cli adm	8	N	N
Slade M, Byford S, Barrett B, Lloyd-Evans B, Gilbert H, Osborn DP, Skinner R, Leese M, Thornicroft G, Johnson S. Br J Psychiatry Suppl. 2010 Aug;53:s14-9. doi: 10.1192/bjp.bp.110.081059. Alternatives to standard acute in-patient care in England: short-term clinical outcomes and cost-effectiveness.	ROM	As a measure in a cost-effectiveness study of different residential facilities	Cli adm	8	N	N
Soergaard KW, Nivison M, Hansen V, Oeiesvold T. BMC Health Serv Res. 2008 May 14;8:103. doi: 10.1186/1472-6963-8-103. Treatment needs and acknowledgement of illness - importance for satisfaction with psychiatric inpatient treatment.	Res	Used as a clinical measure in a satisfaction survey	cli	9	N	N
Sørgaard KW, Rezvy G, Bugdanov A, Sørli T, Bratlid T. Int J Ment Health Syst. 2013 Jan 14;7(1):4. doi: 10.1186/1752-4458-7-4. Treatment needs, diagnoses and use of services for acutely admitted psychiatric patients in northwest Russia and northern Norway.	Res	Comparison of patient groups in 2 different countries	unkno wn	9	N	N
Staring AB, Blaauw E, Mulder CL. Community Ment Health J. 2012 Apr;48(2):150-2. doi: 10.1007/s10597-011-9406-9. Epub 2011 May 10.	ROM	Combining ROM and police data to assess progress	cli	8	N	N

The effects of assertive community treatment including integrated dual diagnosis treatment on nuisance acts and crimes in dual-diagnosis patients		On closer examination of the paper, not actually – it was really an evaluation of the addition of a new treatment component (IDDT) into an ACT programme				
Stephenson LA, Macdonald AJD, Seneviratne G, Waites F, Pawlby S. BJPsych Open. 2018 Apr 19;4(3):119-125. doi: 10.1192/bjo.2018.7. eCollection 2018 May. Mother and Baby Units matter: improved outcomes for both.	Cli	Correlating HoNOS scores with a maternal care index justification of service	Cli adm	8	N	N
Stevens J. A unique day program for adolescents and young adults with eating disorders, program details and outcomes of a 4 year pilot project. <i>EmbaseAsia-Pacific Psychiatry. Conference: 14th Pacific Rim College of Psychiatrists Scientific Meeting. Brisbane, QLD Australia. Conference Publication: (var.pagings). 2 (3) (pp A3), 2010. Date of Publication: October 2010</i>	ROM	Evaluation of eating disorders programme	Cli adm	8	N	N
Stiekema APM, Looijmans A, van der Meer L, Bruggeman R, Schoevers RA, Corpeleijn E, Jörg F. Schizophr Res. 2018 Mar 1. pii: S0920-9964(18)30133-6. doi: 10.1016/j.schres.2018.02.053. [Epub ahead of print] Effects of a lifestyle intervention on psychosocial well-being of severe mentally ill residential patients: ELIPS, a cluster randomized controlled pragmatic trial.	Res	Evaluating a lifestyle intervention programme	cli	8	N	N
Stirone, V; Vigorelli, M; Conte, M; Erlicher, A; Lora, A. Outcome assessment for different care pathway in mental health services. [Italian]. La valutazione dell'Efficacia dei percorsi di cura nei Servizi di Salute Mentale. <i>PsycINFO Ricerca in Psicoterapia. Vol.11(2), 2008, pp. 189-205</i>	Res	HoNOS scores demonstrated care pathways associated with severity of illness	Adm cli	7	N	N
Sugarman, Philip; Walker, Lorraine; Dickens, Geoff. Managing outcome performance in mental health using HoNOS: Experience at St Andrew's Healthcare. <i>PsycINFO Psychiatric Bulletin. Vol.33(8), 2009, pp. 285-288</i>	ROM	Field trial – their conclusion – roms can demonstrate clinical effectiveness – deduced from improving scores paralleling (clinical?) expectations	Cli adm	7	N	N
Sytema S, Wunderink L, Bloemers W, Roorda L, Wiersma D. Acta Psychiatr Scand. 2007 Aug;116(2):105-12. Assertive community treatment in the Netherlands: a randomized controlled trial.	ROM	Evaluation of ACT vs standard community care	Adm cli	8	N	N
Tamasan S., Reisz D., Vasilian C.C. Cognitive impairment and altered emotional status related to type II diabetes mellitus. <i>EmbaseEuropean Psychiatry. Conference: 19th European Congress of Psychiatry, EPA 2011. Vienna Austria. Conference Publication: (var.pagings). 26 (SUPPL. 1) (no pagination), 2011. Date of Publication: March 2011</i>	Res	Survey of diabetics	cli	7	N	N
Taylor, Helen; Stuttford, Maria; Vostanis, Panos. Short-term outcome of young homeless people in contact with a designated mental health service.	Res	Clinical measure in evaluation of the population	Cli adm	9	N	N

PsycINFOThe European Journal of Psychiatry. Vol.21(4), 2007, pp. 268-278.	? ROM ?					
Taylor, John R; Wilkinson, Greg. HoNOS v. GP opinion in a shifted out-patient clinic. PsycINFOPsychiatric Bulletin. Vol.21(8), 1997, pp. 483-485	Res	Weak association between GP ratings and HoNOS scores over time	Cli adm	7	N	N
Theuma, Megan; Read, John; Moskowitz, Andrew; Stewart, Alistair. Evaluation of a New Zealand early intervention service for psychosis. PsycINFONew Zealand Journal of Psychology. Vol.36(3), 2007, pp. 136-145.	ROM	What the title says	adm	8	N	N
Tomasi R, de Girolamo G, Santone G, Picardi A, Micciolo R, Semisa D, Fava S; Gruppo PROGRES. Epidemiol Psichiatri Soc. 2005 Apr-Jun;14(2):77-90. [Drug prescription in Italian Residential Facilities]. [Article in Italian]	Res	HoNOS used without any obvious relevance to the results or conclusions	cli	9	N	N
Tompsett T, Masters K, Donyai P. Neuropsychiatr Dis Treat. 2018 Mar 2;14:681-691. doi: 10.2147/NDT.S151174. eCollection 2018. The utility of novel outcome measures in a naturalistic evaluation of schizophrenia treatment.	Cli ROM	HoNOS used as comparator against the Mental Health Clustering Tool in a study of depot paliperidone	adm	9	N	N
Torrecilla R., Baez Amuedo M.N., Fernandez Monge M. Study of efficiency, physical health and tolerance of paliperidone palmitate in patients with severe mental disorder. EmbaseEuropean Neuropsychopharmacology. Conference: 28th European College of Neuropsychopharmacology, ECNP Congress. Amsterdam Netherlands. Conference Publication: (var.pagings). 25 (SUPPL. 2) (pp S497-S498), 2015. Date of Publication: September 2015	res	Drug trial	cli	8	N	N
Torrecilla Olavarrieta R., Jaen Moreno M.J., Villalobos Martinez E. Study of efficiency and tolerance after changing from olanzapine and risperidone to oral paliperidone in a sample of patients with schizophrenia. EmbaseEuropean Neuropsychopharmacology. Conference: 28th European College of Neuropsychopharmacology, ECNP Congress. Amsterdam Netherlands. Conference Publication: (var.pagings). 25 (SUPPL. 2) (pp S496-S497), 2015. Date of Publication: September 2015	res	Drug trial	cli	8	N	N
Trauer T. Australas Psychiatry. 2004 Jun;12(2):139-44. Use of a routine outcome measure in a consultation-liaison mental health service.	ROM	Justifying the need for CL services – HoNOS used as a clinical measure of severity	adm	8	N	N
Trauer, Tom; Callaly, Tom. Concordance between patients and their case managers using the Health of the Nation Outcome Scales (HoNOS). PsycINFOAustralasian Psychiatry. Vol.10(1), 2002, pp. 24-28	res	Concludes that difference between clinician and patient ratings suggests that they measure different things	cli	9	N	N

Trauer T, Callaly T, Hantz P, Little J, Shields R, Smith J. Br J Psychiatry. 1999 May;174:380-8. Health of the Nation Outcome Scales. Results of the Victorian field trial.	res	What the title says	Adm cli	9	N	N
Trauer T, Callaly T, Hantz P. Aust N Z J Psychiatry. 1999 Jun;33(3):379-84. The measurement of improvement during hospitalisation for acute psychiatric illness.	res	Used as a clinical measure in a study of two psychiatric populations	cli	9	N	N
Trauer, Tom; Nagel, Tricia. Outcome measurement in adult Indigenous mental health consumers. <i>PsycINFOAdvances in Mental Health. Vol.11(1), 2012, pp. 106-116</i>	res	Field trial	Cli	9	N	N
Tsiachristas A, Thomas T, Leal J, Lennox BR. BMJ Open. 2016 Oct 20;6(10):e012611. doi: 10.1136/bmjopen-2016-012611. Economic impact of early intervention in psychosis services: results from a longitudinal retrospective controlled study in England.	ROM	Use of HoNOS and other measures to evaluate EIP	adm	8	N	N
Tulloch, A. D; David, A. S; Thornicroft, G. Exploring the predictors of early readmission to psychiatric hospital. <i>PsycINFOEpidemiology and Psychiatric Sciences. Vol.25(2), 2016, pp. 181-193.</i>	ROM	Clinical measurement in a readmission prediction study	cli	7	N	N
Tulloch AD, Fearon P, David AS. Soc Psychiatry Psychiatr Epidemiol. 2012 Jul;47(7):1181-91. doi: 10.1007/s00127-011-0414-4. Epub 2011 Jul 14. Timing, prevalence, determinants and outcomes of homelessness among patients admitted to acute psychiatric wards.	ROM	A study of homelessness in acute psychiatric inpatients	cli	7	N	N
Tulloch AD, Khondoker MR, Fearon P, David AS. BMC Psychiatry. 2012 Aug 21;12:121. doi: 10.1186/1471-244X-12-121. Associations of homelessness and residential mobility with length of stay after acute psychiatric admission.		HoNOS used as an independent variable in a study of factors influencing LOS	adm	9	N	N
Tulloch AD, Khondoker MR, Thornicroft G, David AS. Epidemiol Psychiatr Sci. 2015 Oct;24(5):402-14. doi: 10.1017/S2045796014000304. Epub 2014 Jun 30. Home treatment teams and facilitated discharge from psychiatric hospital.	ROM	Evaluation of the 'facilitated discharge' function of home treatment teams	cli	8	N	N
Turner, M. A; Boden, J. M; Smith-Hamel, C; Mulder, R. T. Outcomes for 236 patients from a 2-year early intervention in psychosis service <i>PsycINFOActa Psychiatrica Scandinavica. Vol.120(2), 2009, pp. 129-137</i>	Res	HoNOS a clinical measure in evaluation of EIPS	Adm cli	8	N	N
Twomey C, Prina AM, Baldwin DS, Das-Munshi J, Kingdon D, Koeser L, Prince MJ, Stewart R, Tulloch AD, Cieza A. PLoS One. 2016 Nov 30;11(11):e0167103. doi: 10.1371/journal.pone.0167103. eCollection 2016. Utility of the Health of the Nation Outcome Scales (HoNOS) in Predicting Mental Health Service Costs for Patients with Common Mental Health Problems: Historical Cohort Study.	ROM	Most HoNOS domains do not predict costs of treatment Actually, the study does not examine the cost of setting up or implementing the ROM or HoNOS.	adm	7	YE S OR NO	N

Turner MA, Boden JM, Smith-Hamel C, Mulder RT. Acta Psychiatr Scand. 2009 Aug;120(2):129-37. doi: 10.1111/j.1600-0447.2009.01386.x. Epub 2009 Mar 31. Outcomes for 236 patients from a 2-year early intervention in psychosis service.	ROM	Evaluation of an EIP service	Adm cli	8	N	N
Underwood, Lisa; McCarthy, Jane; Tsakanikos, Elias; Howlin, Patricia; Bouras, Nick; Craig, Tom K. J. Health and social functioning of adults with intellectual disability and autism. <i>PsycINFO Journal of Policy and Practice in Intellectual Disabilities</i> . Vol.9(2), 2012, pp. 147-150	ROM	HoNOS as a clinical measure in ID and ASD	Cli adm	9	N	N
Uzoechina, Chinedu E; Hilvering, Pieter; Rogers, Cathryn; O'Brien, Sinead; Pullela, Ananth; Yousafzai, Mushtaq; Meagher, David J. Off-label prescribing in people with recurrent depressive disorder attending a community mental health service. <i>PsycINFO The Psychiatrist</i> . Vol.35(3), 2011, pp. 84-89.	Res ? ROM ?	Clinical measure used as independent variable in examining prescribing practice	cli	9	N	N
Valdes-Stauber J, Kämmerle H, Bachthaler S. Psychother Psychosom Med Psychol. 2015 Dec;65(12):467-75. doi: 10.1055/s-0035-1565087. Epub 2015 Dec 1. [Are there Associations between Embitterment and Clinical as well as Personality Variables by Psychosomatically Ill Inpatients]. [Article in German]	Cli ? ROM ?	Use of HoNOS to investigate bitterness in psychosomatic patients	cli	9	N	N
Valdes-Stauber J, Vietz J, Kilian R. Psychiatr Prax. 2015 May;42(4):191-6. doi: 10.1055/s-0034-1369898. Epub 2014 May 23. [Stepwise outpatient care]. [Article in German]	Res	Using HoNOS scores to justify the presence of an OP psychiatry clinic	adm	7	N	N
Van Duin D, Franx G, Van Wijngaarden B, Van Der Gaag M, Van Weeghel J, Slooff C, Wensing M. Int J Qual Health Care. 2013 Dec;25(6):626-32. doi: 10.1093/intqhc/mzt072. Epub 2013 Oct 31. Bridging the science-to-service gap in schizophrenia care in the Netherlands: the Schizophrenia Quality Improvement Collaborative.	ROM	conclusion of a positive effect from slight HoNOS score improvement in large-scale implementation programme of national guidelines for schizophrenia	Cli adm	8	N	N
van Vugt MD, Kroon H, Delespaul PA, Mulder CL. Psychiatr Serv. 2012;63(5):477-81. doi: 10.1176/appi.ps.201000549. Consumer-providers in assertive community treatment programs: associations with client outcomes.		A study of the effectiveness of consumer providers	Cli adm	8	N	N
van Vugt MD, Kroon H, Delespaul PA, Dreef FG, Nugter A, Roosenschoon BJ, van Weeghel J, Zoeteman JB, Mulder CL. Can J Psychiatry. 2011 Mar;56(3):154-60. Assertive community treatment in the Netherlands: outcome and model fidelity.	ROM res	Outcomes measure comparing different ACT services	cli	8	N	N
Vargas M.L., Quintanilla P., Bega B., Lozano A., Martinez Sanchez J.M., Lopez Gomez D., Martinez De Benito R., Fernandez Atienza C., Lorente Williams M., De La Calle Sanz A. Identification of psychosocial necessities associated with aging in schizophrenia. <i>Embase European Archives of</i>	Res	Study of the needs of ageing in schiz	cli	7	N	N

<i>Psychiatry and Clinical Neuroscience. Conference: 3rd European Conference on Schizophrenia Research: Facts and Visions. Berlin Germany. Conference Publication: (var.pagings). 261 (SUPPL. 1) (pp S100), 2011. Date of Publication: September 2011.</i>						
Vazquez Bourgon J., Hoyuela Zaton F., Gomez-Ruiz E., Cortazar Lopez E., Agueros Perez B., Cuetara Caso J., Gutierrez Ajenjo M.J., Alvaredo Rodriguez C., Rodriguez-Rodriguez P., Pelayo Reventun P., Crespo-Facorro B. Partial psychiatric hospitalization and differences in clinical outcome. <i>EmbaseEuropean Psychiatry. Conference: 24th European Congress of Psychiatry, EPA 2016. Madrid Spain. Conference Publication: (var.pagings). 33 (SUPPL.) (pp S195), 2016. Date of Publication: March 2016</i>	Res	Part of evaluation of a partial hospital	Adm cli	7	N	N
Veerbeek MA, Oude Voshaar RC, Pot AM. Int Psychogeriatr. 2014 Apr 23:1-10. [Epub ahead of print] Effectiveness and predictors of outcome in routine out-patient mental health care for older adults.	Res	people in contact with a psychogeriatric service improve	adm	7	N	N
Veerbeek MA, Oude Voshaar RC, Pot AM. Int J Nurs Stud. 2013 Dec;50(12):1711-9. doi: 10.1016/j.ijnurstu.2013.05.004. Epub 2013 Jun 14. Psychometric properties of the Dutch version of the Health of the Nation Outcome Scales for older adults (HoNOS 65+) in daily care.	ROM	Validation study, looking at the ability of HoNOS 65+ to discriminate or predict	cli	9	N	N
Veerman S., Schulte P., Deijen J.B., De Haan L. Adjunctive memantine in clozapine-treated refractory schizophrenia: A one-year extension study. <i>EmbaseEuropean Psychiatry. Conference: 24th European Congress of Psychiatry, EPA 2016. Madrid Spain. Conference Publication: (var.pagings). 33 (SUPPL.) (pp S108-S109), 2016. Date of Publication: March 2016</i>	Res	Clinical measure in drug trial	cli	8	N	N
Vita A., De Peri L., Barlati S., Cacciani P., Deste G., Poli R., Agrimi E., Cesana B., Sachetti E. Predictors of cognitive improvement and normalization under cognitive remediation in patients with schizophrenia. <i>EmbaseSchizophrenia Research. Conference: 3rd Biennial Schizophrenia International Research Conference. Florence Italy. Conference Publication: (var.pagings). 136 (SUPPL. 1) (pp S76), 2012. Date of Publication: April 2012</i>	Res	Evaluation of cognitive remediation	cli	8	N	N
Ward G, Perera G, Stewart R. Int J Geriatr Psychiatry. 2015 Jun;30(6):639-46. doi: 10.1002/gps.4195. Epub 2014 Oct 14. Predictors of mortality for people aged over 65 years receiving mental health care for delirium in a South London Mental Health Trust, UK: a retrospective survival analysis.	ROM	HoNOS used as an independent variable in a mortality study	cli	9	N	N
Wimalaratne I. An evaluation of a general hospital-based consultation-liaison psychiatry service. <i>EmbaseAustralian and New Zealand Journal of</i>	ROM	Survey of referrals to CL in a general hospital	adm	7	N	N

Psychiatry. Conference: Royal Australian and New Zealand College of Psychiatrists, RANZCP Annual Congress 2013. Sydney Australia. Conference Publication: (var.pagings). 47 (SUPPL. 1) (pp 92), 2013. Date of Publication: May 2013						
Young P. Measuring and thinking about the mental health harms of immigration detention. <i>EmbaseAustralian and New Zealand Journal of Psychiatry. Conference: Royal Australian and New Zealand College of Psychiatrists, RANZCP Annual Congress 2015. South Brisbane, QLD Australia. Conference Publication: (var.pagings). 49 (SUPPL. 1) (pp 84-85), 2015. Date of Publication: May 2015.</i>	Res	Presentation of data of immigrants in detention	Cli adm	9	N	N
Zallo E., Uriarte J.J., Ballesteros J., Zumarraga M. Sociodemographical and clinical characteristics as determinant factors in the derivation to long stay units. <i>EmbaseEuropean Psychiatry. Conference: 18th European Congress of Psychiatry. Munich Germany. Conference Publication: (var.pagings). 25 (SUPPL. 1) (no pagination), 2010. Date of Publication: 2010.</i>	Res	Comparison of patients allocated to , and not allocated to, long-stay units	Cli adm	7	N	N