

Supplementary Table 1 Number of subjects per analysis set in MAGELLAN and MAGELLAN Subpopulation

| Analysis set | MAGELLAN | | | MAGELLAN Subpopulation | | |
|--------------|-------------|------------|-------|------------------------|------------|-------|
| | Rivaroxaban | Enoxaparin | Total | Rivaroxaban | Enoxaparin | Total |
| Randomized | 4050 | 4051 | 8101 | --- | --- | --- |
| Safety | 3997 | 4001 | 7998 | 3218 | 3229 | 6447 |
| mITT Day 35 | 2967 | 3057 | 6024 | 2419 | 2506 | 4925 |
| PP Day 10 | 2938 | 2993 | 5931 | 2385 | 2433 | 4818 |
| mITT Day 10 | 3232 | 3271 | 6503 | 2608 | 2659 | 5267 |
| PP Day 35 | 2516 | 2586 | 5102 | 2066 | 2134 | 4200 |

mITT, modified Intent to Treat; PP, per protocol.

Supplementary Table 2 Reasons to exclude from the MAGELLAN Subpopulation (safety analysis set)

| Exclusion criteria | Rivaroxaban | Enoxaparin/ Placebo | Total |
|--|-------------|------------------------|-------------|
| | N = 779 | N = 772 | N = 1551 |
| Active cancer at randomization | 294 (37.7%) | 290 (37.6%) | 584 (37.7%) |
| Dual antiplatelet therapy at baseline | 246 (31.6%) | 233 (30.2%) | 479 (30.9%) |
| Baseline bronchiectasis/ pulmonary cavitation | 60 (7.7%) | 61 (7.9%) | 121 (7.8%) |
| Active gastroduodenal ulcer | 116 (14.9%) | 116 (15.0%) | 232 (15.0%) |
| Bleeding in 3 months prior to randomization | 126 (16.2%) | 133 (17.2%) | 259 (16.7%) |

Patients who met high risk bleeding exclusion criteria.

Supplementary Table 3 Fatal Bleeding in MAGELLAN

| Treatment Group | Duration of Study Drug | On-set of AE Study Day | Death Study Day | Age/Sex | Relevant Medical History | Bleeding Site |
|--------------------------|------------------------|------------------------|-----------------|------------|---|------------------|
| Rivaroxaban | 2 | 3 | 3 | 62, Male | Bronchiectasis, COPD, septicemia, tuberculosis | Pulmonary |
| Rivaroxaban | 8 | 8 | 9 | 52, Male | Bronchiectasis, COPD, hemoptysis, HF, kidney injury | Pulmonary |
| Rivaroxaban | 11 | 12 | 13 | 44, Female | Active lung cancer, COPD | Pulmonary |
| Rivaroxaban | 3 | 4 | 6 | 79, Male | Bronchiectasis, COPD, pneumonia | Gastrointestinal |
| Rivaroxaban ^a | 5 | 5 | 8 | 70, Male | Afib, COPD, HF, HTN, MI, Renal insufficiency | Intracranial |
| Rivaroxaban ^a | 33 | 33 | 34 | 85, Male | COPD, DM, HTN, pneumonia | Intracranial |
| Rivaroxaban ^a | 11 | 11 | 22 | 70, Male | DM, HTN, liver steatosis | Retroperitoneal |
| Enoxaparin ^a | 2 | 3 | 3 | 86, Male | Suspected bronchial neoplasm, HTN, HF, UTI | Tracheal |

Afib, Atrial Fibrillation; COPD, chronic obstructive pulmonary disease; DM, Diabetes Mellitus; HTN, arterial hypertension; HF, Heart Failure; MI, myocardial infarction; UTI, Urinary Tract infection. ^aFatal Bleeding in MAGELLAN subpopulation.