Health care resource use: anti-depressant use/prescription, NHS outpatient or community mental health team clinics for mental health problems and primary care consultations (Bristol centre only)

Outcome	Self-reported data	Electronic medical	Concordance		
	from	records	between sources		
	questionnaires	N=35	N=35		
	N=35				
Anti-depressant medication	66% (n=23, 0%	43% (n=15, 14%	69% (n=24, 0%		
taken/prescribed	missing)	missing)	missing both		
			sources)		
Number of different anti-depressant taken/prescribed					
1	74% (17/23)	93% (14/15)	50% (12/24)		
2	13% (3/23)	7% (1/15)	0		
3	13% (3/23)	0	0		
NHS outpatient or	11% (n=4, 9%	49% (n=17, 11%	46% (n=16, 0%		
community mental health	missing)	missing)	missing both		
team clinics for mental			sources)		
health problems					
Number of visits					
1	0	18% (3/17)			
2	50% (2/4)	0			
3	0	29% (5/17)			
4	25% (1/4)	0			
7	0	6% (1/17)			
8	0	6% (1/17)			

11	0	6% (1/17)			
12	0	6% (1/17)			
17	0	6% (1/17)			
23	0	6% (1/17)			
24	0	6% (1/17)			
27	0	6% (1/17)			
31	0	6% (1/17)			
Missing	25% (1/4)	0			
At least one GP visit at	46% (n=16, 9%	26% (n=9, 14%	37% (n=13, 3%		
practice for any medical	missing)	missing)	missing both		
condition			sources)		
Number of visits					
0	46% (16/35)	60% (21/35)	31% (11/35)		
1	14% (5/35)	9% (3/35)	3% (1/35)		
2	11% (4/35)	6% (2/35)	0		
3	6% (2/35)	0	0		
4	3% (1/35)	3% (1/35)	0		
5	3% (1/35)	9% (3/35)	3% (1/35)		
6	3% (1/35)	0	0		
7	3% (1/35)	0	0		
9	3% (1/35)	0	0		
Missing	9% (3/35)	14% (5/35)	3% (1/35)		

At least one GP home visit	3% (n=1, 9%	0% (n=0, 14%	77% (n=27, 3%
for any medical condition	missing)	missing)	missing both
			sources)
Number of visits			
0	89% (31/35)	86% (30/35)	60% (42/35)
1	3% (1/35)	0	
Missing	9% (3/70)	14% (5/35)	3% (1/35)