

Supplementary materials

Supplementary Table 1. Clinical and treatment variables based on the use of ticagrelor versus clopidogrel or prasugrel

	Ticagrelor	Clopidogrel or Prasugrel	<i>P</i> -value
Pre-PCI treatment, n/N (%)			
Heparin	98/290 (33.8)	117/240 (48.8)	<.001
Bivalirudin	18/292 (6.2)	32/240 (13.3)	.005
GP IIb/IIIa inhibitors	49/262 (18.7)	55/201 (27.4)	.027
ST resolution, n/N (%)	191/253 (75.5)	119/188 (63.3)	.006
Pre-PCI TIMI score, n/N (%)			.377
TIMI 0	182/292 (62.3)	148/221 (67.0)	
TIMI 1	72/292 (24.7)	41/221 (18.6)	
TIMI 2	24/292 (8.2)	18/221 (8.1)	
TIMI 3	14/292 (4.8)	14/221 (6.3)	
Post-PCI TIMI score, n/N (%)			<.001
TIMI 0	5/263 (1.9)	13/201 (6.5)	
TIMI 1	3/263 (1.1)	2/201 (1.0)	
TIMI 2	6/263 (2.3)	24/201 (11.9)	
TIMI 3	249/263 (94.7)	162/201 (80.6)	
Killip class, n/N (%)			.670
I	214/292 (73.3)	166/240 (69.2)	
II	33/292 (11.3)	33/240 (13.8)	
III	19/292 (6.5)	20/240 (8.3)	

IV	26/292 (8.9)	21/240 (8.8)	
Pre-PCI cardiac arrest, n/N (%)	38/292 (13.0)	18/240 (7.5)	.039
Culprit artery, n/N (%)			0.165
Left main	10/261 (3.8)	5/197 (2.5)	
Left anterior descending	122/261 (46.7)	85/197 (43.1)	
Circumflex	42/261 (16.1)	26/197 (13.2)	
Right coronary	76/261 (29.1)	77/197 (39.1)	
Bypass	11/261 (4.2)	4/197 (2.0)	
Diabetes, n/N (%)	54/290 (18.6)	56/235 (23.8)	.145
Dyslipidemia, n/N (%)	105/289 (36.3)	80/233 (34.3)	.635
Hypertension, n/N (%)	177/292 (60.6)	171/240 (71.3)	.010
Smoker, n/N(%)	135/292 (46.2)	95/237 (40.1)	.156
Cardiovascular disease, n/N (%)	26/290 (9.0)	30/235 (12.8)	.161
Number of vessels with significant disease, n/N(%)			.440
0	2/284	4/205	
1	105/284	70/205	
2	81/284	67/205	
3	96/284	64/205	

Abbreviations: GP, glycoprotein; PCI, percutaneous coronary intervention; TIMI, Thrombolysis in Myocardial Infarction.

Supplementary Table 2. Analysis of 90-day mortality with ticagrelor versus no ticagrelor by propensity score quintile

	Hazard ratio	95% CI	<i>P</i>-value
Ticagrelor (no vs yes)	2.425	1.330–4.421	.004
Propensity score group			
1 st vs 5 th quintile	1.945	.872–4.337	.104
2 nd vs 5 th quintile	1.025	.427–2.461	.956
3 rd vs 5 th quintile	.591	.210–1.665	.320
4 th vs 5 th quintile	.486	.162–1.453	.196

Supplementary Figure 1. Kaplan–Meier curves of (A) 10-day and (B) 30-day cumulative survival with ticagrelor (n = 219) versus no ticagrelor (n = 200), based on propensity score-matched data.

