

Supplementary Table 1. Norms for the Multidimensional Schizotypy Subscales

	<u>Mean</u>	<u>SD</u>
MSS Positive Schizotypy	3.587	4.384
MSS Negative Schizotypy	3.604	4.436
MSS Disorganized Schizotypy	4.008	5.751

Norms based upon 9,366 adults assessed at four universities and on Amazon Mechanical Turk

Supplementary Table 2. Linear Regressions Examining Prediction by the Multidimensional Schizotypy Scale Subscales ($n = 177$)

Criteria:	MSS Positive Schizotypy				MSS Negative Schizotypy				MSS Disorganized Schizotypy			
	B	SE	95%CI		B	SE	95%CI		B	SE	95%CI	
			Lower	Upper			Lower	Upper			Lower	Upper
Global Functioning	-.332**	.127	-.585	-.085	-1.055***	.130	-1.290	-.772	-.665***	.097	-.863	-.478
SIPS-Positive Symptoms	.416***	.044	.325	.501	.141***	.047	.055	.238	.039	.034	-.028	.105
SIPS-Disorganized Symptoms	.092**	.031	.032	.153	.067*	.032	.005	.133	.229***	.026	.178	.278
NSM Negative Symptoms	-.078	.042	-.162	.007	.481***	.058	.356	.587	.150**	.043	.071	.238
NSM Attentional Deficits	.019	.014	-.010	.047	.004	.013	-.020	.030	.088***	.011	.065	.109
Schizotypal Symptoms	.211***	.029	.153	.268	.172***	.040	.094	.255	.027	.025	-.024	.075
Schizoid Symptoms	-.061*	.029	-.116	-.004	.272***	.048	.165	.354	-.009	.021	-.049	.032
Paranoid Symptoms	.075**	.026	.029	.128	.077**	.025	.029	.126	.005	.015	-.025	.036
Alcohol Use	.123	.101	-.068	.334	-.338***	.096	-.531	-.152	.057	.077	-.092	.207
Alcohol Impairment	.018	.011	-.003	.041	-.025**	.009	-.044	-.009	.010	.008	-.006	.025
Drug Use	.035	.072	-.088	.197	-.026	.050	-.125	.076	.058	.051	-.044	.161
Drug Impairment	.002	.015	-.027	.034	-.033**	.010	-.053	-.013	.030**	.011	.008	.050

* $p < .05$ ** $p < .01$ *** $p < .001$

Note: bootstrapped raw regression (B) coefficients with 1000 samples with 95% Confidence Intervals

Each row represents a separate regression analysis in which the three MSS subscales were entered simultaneously as predictors to examine their unique prediction of each of the interview measures.

Supplementary Table 3. Linear Regressions Examining Prediction of Paranoid Personality Disorder Traits by the Multidimensional Schizotypy Scale Subscales ($n = 177$)

MSS Positive Schizotypy					MSS Negative Schizotypy				MSS Disorganized Schizotypy				
<u>Paranoid PD Criteria:</u>	<u>β</u>	<u>B (SE)</u>	<u>95%CI</u>		<u>β</u>	<u>B (SE)</u>	<u>95%CI</u>		<u>β</u>	<u>B (SE)</u>	<u>95%CI</u>		<u>Total R²</u>
			<u>Lower</u>	<u>Upper</u>			<u>Lower</u>	<u>Upper</u>			<u>Lower</u>	<u>Upper</u>	
Reluctance to confide	.204**	.018 (.008)	.004	.035	.242**	.018 (.009)	.002	.035	-.062	-.004 (.004)	-.012	.004	.099***
Doubts loyalty/trustworthiness	.165*	.019 (.010)	.001	.037	.237**	.023 (.010)	.003	.044	.012	.001 (.006)	-.011	.014	.088**
Bears grudges	.009	.001 (.007)	-.012	.014	.139	.011 (.008)	-.003	.029	.096	.006 (.005)	-.003	.017	.032
Perceives attacks and reacts	.068	.004 (.005)	-.004	.014	.050	.003 (.005)	-.006	.007	-.001	.000 (.003)	-.006	.007	.007
Reads hidden meanings	.148	.009 (.007)	-.002	.024	.029	.001 (.003)	-.004	.008	.056	.002 (.003)	-.003	.010	.030
Expects exploitation	.168*	.013 (.007)	.000	.026	.233**	.015 (.008)	.000	.031	.005	.000 (.004)	-.008	.010	.086
Suspects infidelity	.192*	.012 (.006)	.002	.024	.120	.006 (.005)	-.002	.016	-.021	-.001 (.003)	-.006	.005	.051*

* $p < .05$ ** $p < .01$ *** $p < .001$

Note: bootstrapped raw regression (B) coefficients with 1000 samples with 95% Confidence Intervals

Each row represents a separate regression analysis in which the three MSS subscales were entered simultaneously as predictors of paranoid personality disorder traits.

Supplementary Table 4. Linear Regressions Examining Prediction of Schizotypal Personality Disorder Traits by the Multidimensional Schizotypy Scale Subscales (n = 177)

MSS Positive Schizotypy					MSS Negative Schizotypy				MSS Disorganized Schizotypy				
Schizotypal PD Criteria:	β	B (SE)	95%CI		β	B (SE)	95%CI		β	B (SE)	95%CI		Total R^2
			Lower	Upper			Lower	Upper			Lower	Upper	
Lack of close friends	.012	.001 (.005)	-.010	.011	.561***	.038 (.008)	.021	.054	-.187**	-.011 (.004)	-.018	-.004	.324***
Excessive social anxiety	.058	.003 (.003)	-.002	.009	.314***	.013 (.008)	-.002	.009	-.078	-.003 (.003)	-.008	.002	.102***
Ideas of reference	.311***	.036 (.010)	.019	.056	.040	.004 (.008)	-.008	.018	.105	.009 (.007)	-.005	.024	.123***
Odd beliefs	.573***	.073 (.010)	.053	.092	.065	.007 (.008)	-.009	.024	.027	.003 (.007)	-.010	.015	.342***
Unusual perceptual exp.	.323***	.033 (.009)	.013	.050	.119	.010 (.007)	-.002	.027	.067	.005 (.006)	-.007	.018	.136***
Odd thought and speech	.215**	.026 (.009)	.008	.043	.322***	.033 (.009)	.014	.052	.206**	.018 (.009)	.000	.035	.230***
Odd behavior	.003	.000 (.005)	-.008	.010	.371***	.020 (.007)	.006	.035	.068	.003 (.004)	-.005	.011	.149***
Inapp./constricted affect	-.015	-.001 (.004)	-.008	.007	.312***	.019 (.008)	.005	.034	-.007	.000 (.004)	-.009	.009	.097***
Suspiciousness	.360***	.040 (.007)	.027	.050	.287***	.028 (.009)	.010	.046	.039	.003 (.006)	-.008	.015	.229***

* $p < .05$ ** $p < .01$ *** $p < .001$

Note: bootstrapped raw regression (B) coefficients with 1000 samples with 95% Confidence Intervals

Each row represents a separate regression analysis in which the three MSS subscales were entered simultaneously as predictors of paranoid personality disorder traits.

Supplementary Table 5. Variance Accounted for by MSS Subscales and MSS Total Score (*n*=177)

<u>Criteria</u>	<u>Variance accounted by MSS Total Score</u>	<u>Variance accounted for by MSS Subscales</u>	<u>% of Subscale variance accounted by MSS Total</u>
Global Functioning	0.402	0.449	89.5%
SIPS-Positive Symptoms	0.255	0.398	64.1%
SIPS-Disorganized Symptoms	0.388	0.459	84.6%
NSM Negative Symptoms	0.264	0.510	51.8%
NSM Attentional Deficits	0.211	0.314	67.1%
Schizotypal Symptoms	0.314	0.413	75.9%
Schizoid Symptoms	0.091	0.436	20.9%
Paranoid Symptoms	0.109	0.155	70.3%
Alcohol Use	0.005	0.080	6.8%
Alcohol Impairment	0.000	0.052	0.7%
Drug Use	0.004	0.010	43.6%
Drug Impairment	0.002	0.078	2.8%