Appendix A

Search Strategy


Upon an initial database search, some of the articles did not refer to older adults specifically and instead discussed the use of bedrails in nursing home settings. We used the search terms "nursing home", "long-term care", "long-term care", "longterm care", "assisted living", and "assisted-living" to include articles examining the use of bedrails for fall prevention that did not explicitly state "older adult" or its related terms. The use of "n3" in the search terms meant that if the words "bed" and "rail" are within three words of each other in an article, the search results would include that article. The use of the truncated term was to allow inclusion of all terms that had "injur-" as a root.

Table A1
Search Terms Used Based on PICO Question

Population Intervention Outcom	
	ne
older adult* senior* elder* aged geriatric* nursing home* long-term care long-term care assisted living assisted-living	

Thank you to Hayden, A., MLIS, MSc, PhD, University of Calgary for providing support in searching the grey literature.

 $\label{eq:appendix B}$ PRISMA diagram for systematic review on bedrail safety



Appendix C

MMAT scores

Study	Type	Criteria		Score
Agashizula, N., & Wu, W. K. (2009). Effects of Potentially Inappropriate Psychoactive Medications on Falls in US Nursing Home Residents. Drugs & aging, 26(10), 853-860. Quantitative Descriptive		4.1 Met		5/5
		4.2	Met	1
	4.3	Met		
		4.4	Met	1
		4.5	Met	
Canadian Institute for Health Information. (2007). Resident Safety: Characteristics associated with falling in Ontario	Quantitative	4.1	Met	4/5
complex continuing care.	Descriptive	4.2	Not Met	1
i l		4.3	Met	4
l l		4.4	Met	-
		4.5	Met	
Canczuti, E., Wagner, L. M., Brush, B. L., Boltz, M., Renz, S., & Talerico, K. A. (2007). Consequences of an	Quantitative Non-	3.1	Not Met	4/5
ntervention to reduce restrictive side rail use in nursing homes. Journal of the American Geriatrics Society, 55(3), 334- 41.	randomized	3.2	Met	
		3.3	Met	
		3.4	Met	
j		3.5	Met	
Enns, E., Bhoutulla, R., Enva, V., Enuclel, K., & Holroyd-Leduc, J. M. (2014). A controlled quality improvement trial to	Quantitative	2.1	Met	4/5
reduce the use of physical restraints in older hospitalized adults. Journal of the American Geriatrics Society, 62(3), 541-	Randomized	2.2	Met	
545.	Controlled	2.3	Met	
i l		2.4	Not Met	
		2.5	Met	
Fonad, E., Burnard, P., & Emami, A. (2008a). Shielding or caging? Healthcare staffs' views on fall risk and protection in	Qualitative	1.1	Met	5/5
Stockholm. International journal of older people nursing, 3(1), 46-54.		1.2	Met	1
		1.3	Met	
		1.4	Met	
		1.5	Met	
Eonad, E., Emami, A., Wahlin, T B. R., Winblad, B., & Sandmark, H. (2009). Falls in somatic and dementia wards at	Quantitative	4.1	Met	3/5
Community Care Units. Scandinavian journal of caring sciences, 23(1), 2-10.	Descriptive	4.2	Not Met	202
Designation of the Community of the Section of the		4.3	Met	i
	4.4	Not Met		
		4.5	Met	1
and, E., Robins Wahlin, T. B. R., Winhlad, B., Emami, A., & Sandmark, H. (2008b). Falls and fall risk among nursing Qua	Quantitative	4.1	Met	3/5
home residents. Journal of clinical nursing, 17(1), 126-134.		4.2	Not Met	
4.3 4.4 4.5		4.3	Met	
	4.4	Not Met	ĺ	
	Met			
Könke, S., Mühlhauser, I., Gerlach, A., Haut, A., Haustert, B., Möhler, R., & Meyer, G. (2012). Effect of a guideline-	ponent intervention on use of physical restraints in nursing homes: a randomized controlled trial. Randomized 2.2 Me Controlled 2.3 Me	Met	5/5	
based multicomponent intervention on use of physical restraints in nursing homes: a randomized controlled trial.		2.2	Met]
ma, 307(20), 2177-2184.		2.3	Met	
		2.4	Met	
		2.5	Met	
López-Soto, P. J., Sandensky, M. H., Sackett-Lundeen, L. L., De Giorgi, A., Rodriguez-Borrego, M. A., Manfredini, R.,	Quantitative	4.1	Not Met	3/5
& Eabhinn, F. (2016). Temporal patterns of in-hospital falls of elderly patients. Nursing research, 65(6), 435-445. Descriptive	Descriptive	4.2	Not Met	
		4.3	Met	
		4.4	Met	
		4.5	Met	
no, H., Lin, M., & Castle, N. (2011). Physical restraint use and falls in nursing homes: a comparison between residents Quantitative		4.1	Met	5/5
with and without dementia. American Journal of Alzheimer's Disease & Other Dementias &, 26(1), 44-50.	Descriptive	4.2	Met]
,		4.3	Met]
,		4.4	Met]
		4.5	Met	
Meyer, G., Köpke, S., Hasstert, B., & Mühlhasser, I. (2009). Restraint use among nursing home residents: Cross-sectional study and prospective cohort study. Journal of Clinical Nursing, 18, 981-990. 601; 10.1111/j.1365- 2702.2008.02460.x 3.1 Randomized 3.2 3.3 3.4 3.5			Not Met	3/5
	Randomized		Met	
			Not Met	
			Met	
	Met			
Terranova, C., Cardin, F., Martella, B., Amato, B., & Militello, C. (2012). Death after an accidental fall of a 101 year old hospitalized patient. Medico-legal implication of falling in geriatrics. BMC surgery, 12(1), S34.	Qualitative	1.1	Met	5/5
		1.2	Met	
	1.3 Met 1.4 Met 1.5 Met			
Geriatrics & Gerontology International, 7(4), 406-408.	1.1	Met	5/5	
		1.2	Met	
		1.3	Met	
,		1.4	Met	
, <u> </u>		1.5	Met	1

Notes

Thank you to Hayden, A., MLIS, MSc, PhD, University of Calgary for providing support in searching the grey literature