The effects of functionalization of carbon nanotubes on toxicological parameters in mice

Elaheh Mohammadi^{1, 2}, Majid Zeinali³, Mohaddeseh Mohammadi-Sardoo², Maryam Iranpour⁴, Behzad Behnam^{* 5, 6}, Ali Mandegary^{* 5, 2}

¹Student Research Committee, School of Pharmacy, Kerman University of Medical Sciences, Kerman, Iran

²Department of Pharmacology and Toxicology, School of Pharmacy, Kerman University of Medical Sciences, Kerman, Iran

³Biotechnology Research Center, Research Institute of Petroleum Industry (RIPI), Tehran, Iran

⁴Pathology and Stem Cell Research Center, Kerman University of Medical Sciences, Kerman, Iran

⁵Pharmaceutics Research Center, Institute of Neuropharmacology, Kerman University of Medical Sciences, Kerman, Iran

⁶Department of Pharmaceutical Biotechnology, Faculty of Pharmacy, Kerman University of Medical Sciences, Kerman, Iran

*Address Correspondence to: Pharmaceutics Research Center, Institute of Neuropharmacology, Kerman University of Medical Sciences, Haft-Bagh Blvd., Kerman 7616911319, Iran. Tel. +983431325315, Email: <u>behnamb@kmu.ac.ir</u>

Or correspondence to : Pharmaceutics Research Center, Institute of Neuropharmacology, Kerman University of Medical Sciences, Haft-Bagh Blvd., Kerman 7616911319, Iran. Tel. +983431325001, Email: <u>alimandegary@kmu.ac.ir</u>



FIGURE 1. Supplementary information: Digital camera images of tissues to visualize the morphology after 36 and 256 days of final dose. **a** lung, **b** liver.



FIGURE 2. Supplementary information: Digital camera images of tissues to visualize the morphology after 36 and 256 days of final dose. **a** spleen, **b** kidney, **c** testis.