

Supplementary table 2: Structure of social networks of severely mentally ill service users according to the length of time since first contact with psychiatry. Morpheus Study. Belgium 2014–2015 (n=380).

	Length of time since first contact with psychiatry			F value
	Less than 5 years	Between 5 and 15 years	More than 15 years	
Size and composition measures				
Support persons (no.) (i.e. network size)	11.2	12.8	12.2	3.1 *
Professionals (no.)	7.0	7.6	7.9	2.4
Mental health services (no.)	5.4	6.4	6.2	4.9
Health services (no.)	0.9	0.8	1.1	2.7
Social services (no.)	0.9	0.8	1.0	0.4
Justice (no.)	0.1	0.1	0.1	1.8
Generic services (no.)	0.3	0.2	0.2	1.2
Non-professionals (no.)	4.2	5.1	4.5	2.4
Family members (no.)	2.9	3.2	2.9	4.0 *
Friends (no.)	1.2	1.4	1.2	1.3
Diversity of professional services (no.)	3.9	3.7	4.0	0.7
Diversity of professional functions (no.)	5.6	5.8	6.0	1.6
Diversity of types of alters ¹ (no.)	3.3	3.3	3.5	1.3
Cohesion measures				
Density (%)	0.2	0.2	0.2	0.7
Fragmentation (%)	0.6	0.6	0.7	0.9
Components (no.)	5.2	5.2	5.6	0.5
Components (excluding isolates) (no.)	1.6	2	1.8	5.2**
Cliques of maximum size (no.)	1.5	1.7	1.7	0.6
Isolates (no.)	3.6	3.3	3.7	0.9

Size of the largest component (no.)	6	6.8	6.4	1.4
Alters in largest component (%)	0.5	0.5	0.5	1
Size of the largest clique (no.)	4.0	4.8	4.4	3.2
Average size of component (no.)	2.9	3.2	2.8	0.8
Degree centralization (%)	0.2	0.2	0.2	0.6
Betweenness centralization (%)	0.1	0.1	0.1	0.3
Average degree (no.)	2.1	2.5	2.3	1.5
Homophily between professionals and non-professionals (ranges from -1 to 1)	-0.3	-0.3	-0.3	0.1
Homophily between types of alters ¹ (ranges from -1 to 1)	0.1	0.3	0.2	1.6

¹ The types of alters used here are: professionals from mental health services, health services, social services, justice, generic services, and non-professionals.

* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$