

Supplementary File

HIV testing among pregnant women with prenatal care in the United States: 2011-2017 National Survey of Family Growth.

Supplementary Table 1: Characteristics of included and excluded subpopulation, 2011-2017 NSFG

Characteristics	Included Population* N (weighted %)	Excluded Population** N (weighted %)	P-value***
Age			0.666
15-24	505 (28.8)	35 (24.4)	
25-34	825 (54.9)	61 (55.4)	
35-44	236 (16.3)	18 (20.3)	
Education			0.532
No high school diploma	289 (14.6)	22 (13.8)	
High school diploma	494 (28.6)	34 (23.6)	
Some college, no bachelor's degree	450 (26.7)	25 (23.6)	
Bachelor's degree or higher	333 (30.2)	33 (39.0)	
Race/ethnicity			0.094
Non-Hispanic white	616 (49.3)	52 (57.7)	
Hispanic	429 (24.0)	31 (21.3)	
Non-Hispanic black	380 (17.0)	17 (6.0)	
Non-Hispanic other or multiple	141 (9.7)	14 (15.0)	
Marital status			0.351
Married	663 (52.3)	51 (59.6)	
Formerly married	180 (9.0)	10 (5.4)	
Never married	723 (38.7)	53 (35.0)	
Poverty level			0.418
Less than 150%	864 (45.6)	62 (49.2)	
150%-299%	372 (25.0)	24 (17.7)	
300% or above	300 (29.3)	28 (33.2)	
Place of residence			0.724
Rural	245 (16.3)	11 (13.3)	
Urban	1321 (83.7)	103 (86.7)	
Birthplace			0.464
Foreign	249 (15.7)	24 (19.7)	
United States	1317 (84.3)	90 (80.3)	
Completed pregnancy			0.016
1	396 (25.9)	42 (41.4)	
2	405 (27.5)	35 (30.0)	
3 or more	765 (46.6)	37 (28.5)	
PMTCT knowledge	N=1563		0.065
No	1060 (68.3)	90 (81.3)	
Yes	503 (31.7)	24 (18.7)	
Health insurance coverage (N=1565)	N=1565		0.350
Yes	1209 (77.4)	85 (82.1)	
No	356 (22.6)	29 (17.9)	
Total	1566 (92.5)	144 (7.5)	

* Women with pregnancy ending within 12 months prior to interview that responded “Yes” or “No” to testing for HIV during prenatal care.

**Women with pregnancy ending within 12 months prior to interview that responded “Never went for prenatal care” or “Don’t know” to testing for HIV during prenatal care.

***Rao-Scott chi-square test. Bold text indicates statistical significance